

The fully residually F quotients of $F * \langle x, y \rangle$

Nicholas W.M. Touikan
 Department of Mathematics and Statistics
 McGill University
 email: touikan@math.mcgill.ca

May 2, 2009

Abstract

We describe the fully residually F groups; or limit groups relative to F ; that arise as the coordinate groups of irreducible systems of equations in two variables over a free group F that have coefficients in F . We use the structure theory of f.g. fully residually free groups and Bass-Serre theory. In particular we use foldings sequences for subgroups of fundamental groups of graphs of groups.

Contents

1	Introduction	2
1.1	F -groups and Algebraic Geometry	3
1.2	Splittings	4
1.2.1	Relative presentations	6
1.3	The cyclic JSJ decomposition	6
1.4	The Structure of $\text{Hom}_F(F_{R(S)}, F)$	9
2	The Classification Theorem	10
2.1	A description of Solutions	13
2.2	Examples and Questions	14
2.2.1	Questions	15
3	Tools for fully residually F groups:	16
3.1	Resolutions	16
3.2	Some notions of complexity	17
3.3	Rational Equivalence	19
4	Bass-Serre Theory Techniques	19
4.1	Induced splittings and $\mathcal{G}(A)$ -graphs	20
4.1.1	Basic definitions	20
4.1.2	Moves on $\mathcal{G}(A)$ -graphs	22
4.1.3	The Folding Process	23
4.1.4	Transmission graphs and 1-acylindrical splittings	24

4.2	When $F_{R(S)}$ splits as a graph of groups with two cycles . .	25
4.3	The Weidmann Nielsen Technique for Groups acting on Trees	27
4.4	Tricks that work for fully residually F groups	27
5	Coordinate Groups that are freely decomposable modulo F	30
6	Splittings with one nonabelian vertex group and at least one abelian vertex group	30
7	Splittings with two or more nonabelian vertex groups	32
7.1	Maximal Abelian Collapses	32
7.1.1	The strategy	32
7.2	The one edge case	33
7.3	The two edge case	40
7.4	The three edge case	46
8	Splittings with one vertex group	48
8.1	First Classifications	48
8.2	The case where $\tilde{F}$ is not freely decomposable modulo F . .	50
8.2.1	Controlling induced splittings	51
8.2.2	Controlling the Strict Quotients	57
8.2.3	The two edge case	58
8.3	The case where $\tilde{F}$ is freely decomposable modulo F	59

1 Introduction

Systems of equations over free groups have been a very important and fruitful subject of study in the field of combinatorial and geometric group theory. A major achievement was the algorithm due to Makanin and Razborov [Mak82, Raz87] which produces a complete description of the solution set of an arbitrary finite system of equations over a free group. Their methods, however, are algorithmic and use surprisingly little algebra.

In [BMR99] Baumslag, Myasnikov and Remeslennikov developed notions of algebraic geometry for groups that applied to CSA groups, which characterized the coordinate groups of irreducible varieties as f.g. fully residually free groups. In their papers [KM98a, KM98b] Kharlampovich and Myasnikov, by improving the techniques of Makanin and Razborov, obtain an algebraic description of f.g. fully residually free groups as subgroups of the Lyndon group $F^{\mathbb{Z}[t]}$ as well as an algebraic description of the set of homomorphisms of an arbitrary finitely presented group into a free group.

Independently, Sela in [Sel01], using the theory of actions on $\mathbb{R}$ -trees (which itself has a very rich genealogy) obtained essentially the same description of the set of homomorphisms of an arbitrary finitely presented group into a free group.

The main players in this algebraic theory are the fully residually free, or limit, groups. Although it is relatively easy to construct many examples of these groups, there are still relatively few systematic classifications. The

theory is so rich that it is hard to say anything significant about the large class of fully residually free groups in terms of their rank, i.e. the minimal number of generators.

The case of equations over two variables however has been the object of much study: [Hme71, Wic72, Ozh83] and in fact provides a wealth of structure while remaining tractable. The purpose of this paper is to describe the fully residually F groups that arise from irreducible systems of equations over F in two variables. Our techniques come from the algebraic structure theorems for fully residually free groups that arose from the works of Kharlampovich, Myasnikov, and Sela on the Tarski problems.

1.1 F -groups and Algebraic Geometry

A complete account of the material in this section can be found in [BMR99]. Fix a free group F . An equation in variables x, y over F is an expression of the form

$$E(x, y) = 1$$

where $E(x, y) = f_1 z_1^{m_1} \dots z_n^{m_n} f_{n+1}$; $f_i \in F, z_j \in \{x, y\}$ and $m_k \in \mathbb{Z}$.

We view an equation as an element of the group $F[x, y] = F * F(x, y)$. A *solution* of an equation is a substitution

$$x \mapsto g_1, y \mapsto g_2; g_i \in F \quad (1)$$

so that in F the product $E(g_1, g_2) =_F 1$. A *system of equations* in variables x, y ; $S(x, y) = S$; is a subset of $F[x, y]$ and a solution of $S(x, y)$ is a substitution as in (1) so that all the elements of $S(x, y)$ vanish in F .

Definition 1.1. A group G equipped with a distinguished monomorphism

$$i : F \hookrightarrow G$$

is called an F -group we denote this (G, i) . Given F -groups (G_1, i_1) and (G_2, i_2) , we define an F -homomorphism to be a homomorphism of groups f such that the following diagram commutes:

$$\begin{array}{ccc} G_1 & \xrightarrow{f} & G_2 \\ i_1 \uparrow & \nearrow i_2 & \\ F & & \end{array}$$

We denote by $\text{Hom}_F(G_1, G_2)$ the set of F -homomorphisms from (G_1, i_1) to (G_2, i_2) .

In the remainder the distinguished monomorphisms will in general be obvious and not explicitly mentioned. It is clear that every mapping of the form (1) induces an F -homomorphism $\phi(g_1, g_2) : F[x, y] \rightarrow F$, it is also clear that every $f \in \text{Hom}_F(F[x, y], F)$ is induced from such a mapping. It follows that we have a natural bijective correspondence

$$\text{Hom}_F(F[x, y], F) \leftrightarrow F \times F = \{(g_1, g_2) | g_i \in F\}$$

Definition 1.2. Let $S = S(x, y)$ be a system of equations. The subset

$$V(S) = \{(g_1, g_2) \in F \times F \mid x \mapsto g_1, y \mapsto g_2 \text{ is a solution of } S\}$$

is called the *algebraic variety* of S .

We have a natural bijective correspondence

$$\text{Hom}_F(F[x, y]/ncl(S), F) \leftrightarrow V(S)$$

Definition 1.3. The *radical* of S is the normal subgroup

$$Rad(S) = \bigcap_{f \in \text{Hom}_F(F[x, y]/ncl(S), F)} \ker(f)$$

and we denote the *coordinate group* of S

$$F_{R(S)} = F[x, y]/Rad(S)$$

It follows that there is a natural bijective correspondence

$$\text{Hom}_F(F[x, y]/ncl(S), F) \leftrightarrow \text{Hom}_F(F_{R(S)}, F)$$

so that $V(S) = V(Rad(S))$. We say that $V(S)$ or S is *reducible* if it is a union

$$V(S) = V(S_1) \cup V(S_2); V(S_1) \subsetneq \cup V(S) \supsetneq V(S_2)$$

of algebraic varieties. An F -group G is said to be *fully residually F* if for every finite subset $P \subset G$ there is some $f_P \in \text{Hom}_F(G, F)$ such that the restriction of f_P to P is injective.

Theorem 1.4 ([BMR99]). *S is irreducible if and only if $F_{R(S)}$ is fully residually F .*

Theorem 1.5 ([BMR99]). *Either $F_{R(S)}$ is fully residually F or*

$$V(S) = V(S_1) \cup \dots \cup V(S_n)$$

where the $V(S_i)$ are irreducible and there are canonical epimorphisms $\pi_i : F_{R(S)} \rightarrow F_{R(S_i)}$ such that each $f \in \text{Hom}_F(F_{R(S)}, F)$ factors through some π_i .

Corollary 1.6. *If $F[x, y]/ncl(S)$ is fully residually F then $F_{R(S)} = F[x, y]/ncl(S)$.*

1.2 Splittings

We assume the reader is familiar with Bass-Serre theory, so we only describe enough to explain our notation.

Definition 1.7. A *graph of groups* $\mathcal{G}(A)$ consists of a connected directed graph A with vertex set VA and edges EA . A is directed in the sense that to each $e \in EA$ there are functions $i : EA \rightarrow VA, t : EA \rightarrow VA$ corresponding to the *initial* and *terminal* vertices of edges. To A we associate the following:

- To each $v \in VA$ we assign a *vertex group* A_v .
- To each $e \in EA$ we assign an *edge group* A_e .

- For each edge $e \in EA$ we have monomorphisms

$$i_e : A_e \rightarrow A_{i(e)}, \quad t_e : A_e \rightarrow A_{t(e)}$$

we call the maps i_e, t_e *boundary monomorphisms* and the images of these maps *boundary subgroups*.

We also formally define the following expressions: for each $e \in EA$

$$(e^{-1})^{-1} = e, \quad i(e^{-1}) = t(e), \quad t(e^{-1}) = i(e), \quad i_{e^{-1}} = t_e, \quad t_{e^{-1}} = i_e$$

A graph of groups has a fundamental group denoted $\pi_1(\mathcal{G}(A))$. We say that a group *splits* as the fundamental group as a graph of groups if $G = \pi_1(\mathcal{G}(A))$ and refer to the data $D = (G, \mathcal{G}(A))$ as a *splitting*.

Definition 1.8. A sequence of the form

$$a_0, e_1^{\epsilon_1}, a_1, e_2^{\epsilon_2}, \dots, e_n^{\epsilon_n}, a_n$$

where $e_1^{\epsilon_1}, \dots, e_n^{\epsilon_n}$ is an edge path of A and where $a_i \in A_{i(e_{i+1}^{\epsilon_{i+1}})} = A_{t(e_1^{\epsilon_1})}$ is called a $\mathcal{G}(A)$ -path.

Definition 1.9. We denote by $\pi_1(G(A), u)$ the group generated by $\mathcal{G}(A)$ paths whose underlying edge path is a loop at u .

We have in particular that $\pi_1(\mathcal{G}(A), u) \approx \pi_1(\mathcal{G}(A))$.

Convention 1.10. If $F_{R(S)}$ is the fundamental group of $\mathcal{G}(A)$, then we will always assume that the basepoint v is the vertex $v \in VA$ such that $F \leq A_v$.

Definition 1.11 (Moves on $\mathcal{G}(A)$). We have the following moves on $\mathcal{G}(A)$ that do not change the fundamental group.

- *Change the orientation of edges* in $\mathcal{G}(A)$, and relabel the boundary monomorphisms.
- *Conjugate boundary monomorphisms*, i.e. replace i_e by $\gamma_g \circ i_e$ where γ_g denotes conjugation by g and $g \in A_{i(e)}$.
- *Slide*, i.e. if there are edges e, f such that $i_e(A_e) = i_f(A_f)$ then we change X by setting $i(f) = t(e)$ and replacing i_f by $t_e \circ i_e^{-1} \circ i_f$.
- *Folding*, i.e. if $i_e(A_e) \leq A \leq A_{i(e)}$, then replace $A_{t(e)}$ by $A_{t(e)} *_{A_e} A$, replace A_e by a copy of A and change the boundary monomorphism accordingly.
- *Collapse an edge* e , i.e. for some edge $e \in EA$, take the subgraph $star(e) = \{i(e), e, t(e)\}$ and consider the quotient of the graph A , subject to the relation $\sim$ that collapses $star(e)$ to a point. The resulting graph $A' = A / \sim$ is again a directed graph. Denote the equivalence class $v' = [star(e)] \in A'$, then we have $A'_{v'} = A_{i(e)} *_{A_e} G_{t(e)}$ or $A_{i(e)} *_{A_e}$ depending whether $i(e) = t(e)$ or not. For each edge f of A incident to either $i(e)$ or $t(e)$, we have boundary monomorphisms $A_f \rightarrow A'_{v'}$ given by $i'_f = j \circ i_f$ or $t'_f = j \circ t_f$, where j is the one of the inclusion $A_{t(e)} \subset A'_{v'}$ or $A_{i(e)} \subset A'_{v'}$.
- *Conjugation*, i.e. for some $g \in G$ replace all the vertex groups A_v by A_v^g and postcompose boundary monomorphisms with γ_g (which denotes conjugation by g).

1.2.1 Relative presentations

Although the use of graphs of groups is critical, they are notationally cumbersome. We therefore recall how to obtain relative presentations from graphs of groups, and explain the graphical notation we will use.

Let $G_1, \dots, G_n$ be groups with presentations $\langle X_1 \mid R_1 \rangle, \dots, \langle X_n \mid R_n \rangle$ (resp.) and $t_1, \dots, t_k$ a set of letters. Let R denote a set of words in $\bigcup X_i^{\pm 1} \cup \{t_1, \dots, t_k\}^{\pm 1}$ then we will define the *relative presentation*

$$\langle G_1, \dots, G_n, t_1, \dots, t_k \mid R \rangle$$

to be the group defined by the presentation

$$\langle X_1, \dots, X_n, t_1, \dots, t_k \mid R_1, \dots, R_n, R \rangle$$

If G is the fundamental group of a graph of groups $\mathcal{G}(A)$ with vertex groups $G_1, \dots, G_n$ and cyclic edge groups we can give G a relative presentation as follows:

- (A) Take a spanning tree T of the underlying graph A .
- (B) For each edge e of T we can assume that $G_{i(e)} \cap G_{t(e)} = G_e$, and therefore can form an iterated amalgam.
- (C) For each edge f not in T we add a “stable letter” t_f and the relation $t_f = (t_f)^{-1} \alpha t_f = \alpha'$; where α and α' are the images in $G_{i(f)}$ and $G_{e(f)}$ resp. of a generator of G_f via boundary monomorphisms.

The resulting presentation gives a group isomorphic to G , although it depends on the choice of spanning tree. We can take the underlying graph A and encode this relative presentation by labelling as follows:

- Vertices are labelled by the vertex groups.
- Edges in the spanning tree T are represented by undirected edges labelled by a generator of the edge group.
- Edges f not in the spanning tree T are directed, labelled by the corresponding stable letter t_f , moreover the endpoints are decorated by the elements α, α' as defined in (C) above.

Throughout this paper we will switch freely between words represented as $G(A)$ -paths and words in generators of relative presentations.

1.3 The cyclic JSJ decomposition

Definition 1.12. An *elementary cyclic splitting* D of G is a splitting of G as either a free product with amalgamation or an HNN extension over a cyclic subgroup. We define the *Dehn twist along* D , δ_D , as follows.

- If $G = A *_{\langle \gamma \rangle} B$ then

$$\delta_D(x) = \begin{cases} x & \text{if } x \in A \\ x^\gamma & \text{if } x \in B \end{cases}$$

- If $G = \langle A, t \mid t^{-1} \gamma t = \beta \rangle, \gamma, \beta \in A$ then

$$\delta_D(x) = \begin{cases} x & \text{if } x \in A \\ t\beta & \text{if } x = t \end{cases}$$

A Dehn twist generates a cyclic subgroup of $\text{Aut}(G)$. A splitting such that all the edge groups are nontrivial and cyclic is called a *cyclic splitting*.

We can generalize the notion of a Dehn twist to arbitrary cyclic splittings.

Definition 1.13. let D be a cyclic splitting of G with underlying graph A and let e be an edge of A . Then a *Dehn twist* along e is an automorphism that can be obtained by collapsing all the other edges in A to get a splitting D' of G with only the edge e and applying one of the applicable automorphisms of Definition 1.12

Definition 1.14. (i) A subgroup $H \leq G$ is elliptic in a splitting D if H is conjugate into a vertex group of D , otherwise we say it is hyperbolic.

(ii) Let D and D' be two elementary cyclic splittings of a group G with boundary subgroups C and C' , respectively. We say that D' is *elliptic in D* if C' is elliptic in D . Otherwise D' is *hyperbolic in D* .

A splitting D of an F -group is said to be *modulo F* if the subgroup F is contained in a vertex group.

The following is proved in [RS97]:

Theorem 1.15. (i) Let G be freely indecomposable (modulo F) and let D', D be two elementary cyclic splittings of G (modulo F). D' is elliptic in D if and only if D is elliptic in D' .

(ii) Moreover if D' is hyperbolic in D then G admits a splitting E such that one of its vertex groups is the fundamental group $Q = \pi_1(S)$ of a punctured surface S such that the boundary subgroups of Q are puncture subgroups. Moreover the cyclic subgroups $\langle d \rangle, \langle d' \rangle$ corresponding to D, D' respectively are both conjugate into Q .

Definition 1.16. A subgroup $Q \leq G$ is a quadratically hanging (QH) subgroup if for some cyclic splitting D of G , Q is a vertex group that arises as in item (ii) of Theorem 1.15.

Not every surface with punctures can yield a QH subgroup. By Theorem 3 of [KM98a], the projective plane with puncture(s) and the Klein bottle with puncture(s) cannot give QH subgroups.

Definition 1.17. (i) A QH subgroup Q of G is a *maximal QH* (MQH) subgroup if for any other QH subgroup Q' of G , if $Q \leq Q'$ then $Q = Q'$.

(ii) Let D be a splitting of G with Q be a QH vertex subgroup and let C be a splitting of Q with boundary subgroup $\langle c \rangle$ then there is a splitting D' of G called a *refinement of D along C* such that D is obtained from a collapse of D' along an edge whose corresponding group is $\langle c \rangle$.

Definition 1.18. (i) A splitting D is *almost reduced* if vertices of valence one and two properly contain the images of edge groups, except vertices between two MQH subgroups that may coincide with one of the edge groups.

(ii) A splitting D of G is *unfolded* if D can not be obtained from another splitting D' via a folding move (See Definition 1.11).

Theorem 1.19 (Proposition 2.15 of [KM05]). *Let H be a freely indecomposable modulo F f.g. fully residually F group. Then there exists an almost reduced unfolded cyclic splitting D called the cyclic JSJ splitting of H modulo F with the following properties:*

- (1) *Every MQH subgroup of H can be conjugated into a vertex group in D ; every QH subgroup of H can be conjugated into one of the MQH subgroups of H ; non-MQH [vertex] subgroups in D are of two types: maximal abelian and non-abelian [rigid], every non-MQH vertex group in D is elliptic in every cyclic splitting of H modulo F .*
- (2) *If an elementary cyclic splitting $H = A *_C B$ or $H = A *_C$ is hyperbolic in another elementary cyclic splitting, then C can be conjugated into some MQH subgroup.*
- (3) *Every elementary cyclic splitting $H = A *_C B$ or $H = A *_C$ modulo F which is elliptic with respect to any other elementary cyclic splitting modulo F of H can be obtained from D by a sequence of moves given in Definition 1.11.*
- (4) *If D_1 is another cyclic splitting of H modulo F that has properties (1)-(2) then D_1 can be obtained from D by a sequence of slidings, conjugations, and modifying boundary monomorphisms by conjugation (see Definition 1.11.)*

Convention 1.20. Unless stated otherwise, instead of saying the *cyclic JSJ decomposition of $F_{R(S)}$ modulo F* , we will simply say the *JSJ of $F_{R(S)}$* .

Definition 1.21. Suppose first that G is freely indecomposable. Given D , a cyclic JSJ decomposition of $F_{R(S)}$ modulo F , we define the group Δ of *canonical F -automorphisms with respect to D* of $F_{R(S)}$ to be generated by the following:

- Dehn twists along edges of D that fix F pointwise;
- automorphisms of the MQH groups that fix edge groups pointwise;
- automorphisms of the abelian vertex groups that fix edge groups point wise.

If G is freely decomposable modulo F then Δ is generated by the extensions of the canonical F -automorphisms of its freely indecomposable free factors.

Convention 1.22. Unless stated otherwise, instead of saying the *canonical F -automorphisms with respect to D* where D is a JSJ, we will simply say the *canonical automorphisms of $F_{R(S)}$* .

The following Theorem is proved in [KM98b].

Theorem 1.23. *If $F_{R(S)} \neq F$ is fully residually free and is freely indecomposable (modulo F) then it admits a non trivial cyclic splitting modulo F .*

Corollary 1.24. *If $F_{R(S)} \neq F$ then it has a nontrivial JSJ.*

1.4 The Structure of $\text{Hom}_F(F_{R(S)}, F)$

Definition 1.25. A *Hom diagram* for $\text{Hom}_F(G, F)$, denoted $\text{Diag}(G, F)$, consists of a finite directed rooted tree T with root v_0 , along with the following data:

- To each vertex, except the root, v of T we associate a fully residually F group $F_{R(S_v)}$.
- The group associated to each leaf of T is a free product $F * F(Y)$ where Y is some set of variables.
- To each edge e with initial vertex v_i and terminal vertex v_t we have a proper F -epimorphism $\pi_e : F_{R(S_{v_i})} \rightarrow F_{R(S_{v_t})}$

We point out that in the work of Sela, the *Hom* diagram is called a *Makanin-Razborov* diagram (relative to F) and that our fully residually F groups are *limit groups* (relative to F). The following theorem gives a finite parametrisation of the solutions of systems of equations over a free group.

Theorem 1.26 ([KM05, Sel01]). *For any system of equations $S(x_1, \dots, x_n)$ there exists a Hom diagram $\text{Diag}(F_{R(S)}, F)$ such that for every $f \in \text{Hom}_F(F_{R(S)}, F)$ there is a path*

$$v_0, e_1, v_1, e_2, \dots, e_{m+1}, v_{m+1}$$

from the root v_0 to a leaf v_{m+1} such that

$$f = \rho \circ \pi_{v_{m+1}} \circ \sigma_{v_m} \circ \dots \circ \sigma_{v_1} \circ \pi_{e_1}$$

where the σ_{v_j} are canonical F -automorphisms of $F_{R(S_{v_j})}$, the π_j are epimorphisms $\pi_j : F_{R(S_{v_j})} \rightarrow F_{R(S_{v_{j+1}})}$ inside $\text{Diag}(F_{R(S)}, F)$, and ρ is any F -homomorphism $\rho : F_{R(S_{v_{m+1}})} \rightarrow F$ from the free group $F_{R(S_{v_{m+1}})}$ to F .

Definition 1.27. Let D be a cyclic splitting of $F_{R(S)}$. If v is a valence 1 vertex of A , the graph underlying D , and A_v is cyclic, then it is called a *hair*.

Definition 1.28. Let D be the JSJ of $F_{R(S)}$ and let D' be splitting of $F_{R(S)}$ obtained by collapsing hairs into the adjacent vertex groups. Then D' is called the *hairless JSJ* of $F_{R(S)}$.

Lemma 1.29. *Theorem 1.26 holds if we replace the canonical automorphisms w.r.t. the JSJ by the canonical automorphisms w.r.t. the hairless JSJ.*

Proof. Since the hairless JSJ is a collapse of the JSJ, we have that the canonical automorphisms w.r.t. the hairless JSJ are a subset of the canonical automorphisms. On the other hand all the Dehn twists associated to hairs are trivial so the sets of automorphisms are equal. $\square$

Convention 1.30. Unless stated otherwise, we will always replace the JSJ by the hairless JSJ.

2 The Classification Theorem

So far the only comprehensive classification theorems of fully residually free groups in terms of the number of generators are the following:

Theorem 2.1. *[FGM⁺98] If G is fully residually free group. Then*

1. *if $\text{Rank}(G) = 1$ then G is infinite cyclic.*
2. *if $\text{Rank}(G) = 2$ then G is free or free abelian of rank 2.*
3. *if $\text{Rank}(G) = 3$ then G is either free or free abelian of rank 3 or, G is isomorphic to a free rank one extension of a centralizer of a free group of rank 2.*

This next result follows from the results of Appel and Lorenc, although their proofs contained gaps. A correct proof was provided by Chiswell and Remeslennikov.

Theorem 2.2. *[App68, Lor68, CR00] If $S(x)$ is an irreducible system of equations over F in one variable then*

$$F_{R(S)} \approx \begin{cases} F \\ F * \langle t \rangle \\ F *_u \text{Ab}(u, t) \end{cases} \quad (2)$$

where $\text{Ab}(u, t)$ denotes the free abelian group with basis $\{u, t\}$.

The class of coordinate groups of irreducible systems of equations in two variables over F is much more varied. The classification we give will be in terms of cyclic JSJ decompositions modulo F . Specifically we will describe the cyclic graphs of groups in terms of the underlying graph, vertex groups, and edge groups. We will also indicate where, up to rational equivalence, the variables x, y are sent.

The groups will be organized as follows: (A) will be the freely decomposable modulo F groups; (B) will be the groups whose JSJ has only one vertex; (C) will be the groups whose JSJ has more than one vertex group, but such that only one of them is nonabelian; (D), (E), and (F) will be the remaining cases. This classification closely follows the proof.

Definition 2.3. If $F_{R(S)}$ has a JSJ D with more than 2 nonabelian vertex groups, then we will call a *cyclic collapse* of D a graph of groups obtained by performing a maximal number of edge collapses (See Definition 1.11) while ensuring that the resulting graph of groups has at least two nonabelian vertex groups.

Definition 2.4. The *first Betti number* of G denoted $b_1(G)$ is the rank of the torsion-free summand of the abelianization of G .

Convention 2.5. Throughout the paper whenever $F_{R(S)}$ is given as the fundamental group of a graph of groups modulo F , we shall denote by $\tilde{F}$ the vertex group that contains F .

Theorem 2.6 (The main Theorem). *Let F be a free group of rank $N \geq 2$. If $S = S(x, y)$ is an irreducible system of equations over F in two variables then $F_{R(S)}$ must be one of the following.*

(A) If $F_{R(S)}$ is freely decomposable then,

$$F_{R(S)} \approx \begin{cases} F * \langle t \rangle \\ F * H; \text{ where } H \text{ is fully residually free of rank 2} \\ (F *_u \text{Ab}(u, t)) * \langle s \rangle \end{cases}$$

(B) If the cyclic JSJ decomposition of $F_{R(S)}$ modulo F has one vertex group and one edge group then, up to rational equivalence, we can arrange to have a relative presentation

$$\begin{array}{c} \overline{y} \\ \curvearrowright \\ \beta \widetilde{F}^{\beta'} \end{array}$$

such that $\widetilde{F} = \langle F, \overline{x}, \beta' \rangle$ and $\beta \in \langle F, \overline{x} \rangle$. If the JSJ has two edges, then, up to rational equivalence, we can arrange to have a relative presentation

$$\left(\begin{array}{c} \alpha \\ \alpha' \widetilde{F}_{\beta'}^{\beta} \end{array} \right)$$

such that $\widetilde{F} = \langle F, \alpha', \beta' \rangle$ and $\alpha \in F$. In both cases the vertex group $\widetilde{F}$ is either $F * \langle z \rangle$ or a cyclic HNN extension of $F * \langle z \rangle$ and we must have $b_1(\widetilde{F}) = N + 1$.

(C) If the cyclic JSJ decomposition of $F_{R(S)}$ modulo F has only the non-abelian vertex group $\widetilde{F}$ and all other vertex groups are abelian. Then the possible graphs of groups are:

$$\mathcal{G}(A) = u \bullet \text{---} \bullet v, v \bullet \text{---} \bullet u \text{---} \bullet w, \text{ or } \begin{array}{c} \curvearrowright \\ u \bullet \text{---} \bullet v \end{array}$$

with $A_u = \widetilde{F}$ and the other vertex groups free abelian. Moreover we have $b_1(\widetilde{F}) \leq N + 1$ if there are two vertex group and $\widetilde{F} = F$ if there are three vertex groups. In the case where we have two edges and two vertices, then exactly one of the boundary subgroups must be conjugate into F itself.

Otherwise, the cyclic JSJ decomposition modulo F of $F_{R(S)}$ is given by a graph of groups $\mathcal{G}(A)$ that has at least two non abelian vertex groups, in which case either:

(D) The JSJ of $F_{R(S)}$ has the cyclic collapse $u \bullet \text{---} \bullet v$. In all cases we have two nonabelian vertex groups. Up to rational equivalence we can arrange to have the following:

(I) The JSJ of $F_{R(S)}$ has one edge and we have the relative presentation $F_{R(S)} = \widetilde{F} *_p H$. Moreover either:

1. $\widetilde{F} \neq F$ then $\widetilde{F} = F *_u \text{Ab}(u, t)$, $p \notin F$ and H is free of rank 2 and generated by $\overline{x}$ and $\overline{y}$; or
2. $\widetilde{F} = F$ and H is free of rank 2 and generated by $\overline{x}, \overline{y}$ and $p \in F$; moreover
3. H is a QH subgroup then we can arrange so that $F_{R(S)} = F *_u \langle x, y, p | [x, y] = p \rangle$.

(II) The JSJ of $F_{R(S)}$ has two edges and either:

1. there are only two vertex groups, then the JSJ is

$$F_{R(S)} = \tilde{F} \underset{u}{\text{---}} H'_{q'} \bigg) \quad \text{where the subgroups } u \text{ and } q \text{ are not conjugate in } F_{R(S)}.$$

The subgroup $H'_{q'} = H$ is free of rank 2 and $\tilde{F}$ and H are as in (I) above; or

2. there are three vertices and we have a presentation

$$F_{R(S)} = F *_u H *_q \text{Ab}(q, t)$$

H is free of rank 2 and the subgroup $H *_u \text{Ab}(q, t)$ is generated by $\overline{x}, \overline{y}$ and $u \in F$. Moreover u and q may be conjugate.

(III) The JSJ of $F_{R(S)}$ has three edge groups, then the only possibility is

$$F_{R(S)} = F \underset{u}{\text{---}} \alpha H^{\alpha'} \underset{p}{\text{---}} \text{Ab}(u, s)$$

moreover α may be conjugate to either u or p , but not both.

(E) The JSJ of $F_{R(S)}$ has the cyclic collapse $u \bullet \text{---} v$ and again there are two non abelian vertex groups. Up to rational equivalence we can arrange to have the following:

(I) The JSJ of $F_{R(S)}$ has two edges and we have a relative presentation:

$$F_{R(S)} = \tilde{F}^\beta \xrightarrow[\alpha]{\overline{y}} \gamma H$$

moreover either:

1. $\tilde{F} = F$ and H is a free group of rank 2 generated by $\overline{x}, \alpha, \gamma$. Where $\gamma = \overline{y}^{-1} \beta \overline{y}$, with $\beta \in F$
2. $\tilde{F} = F *_u \text{Ab}(u, s)$ and H is a free group of rank 2 generated by $\overline{x}$ and $\alpha \in F$.

(II) The JSJ of $F_{R(S)}$ has three edges and either:

1. We have the relative presentation

$$F_{R(S)} \approx F^\gamma \xrightarrow[\alpha]{\overline{y}} \gamma' H'_{q'} \bigg)$$

where $H = \gamma' H'_{q'}$ is free of rank 2, moreover α and γ are not conjugate, but it is possible for q to be conjugate to one of them. The vertex groups are generated as in (II).1. above.

2. $F_{R(S)} = F^\beta \xrightarrow[\alpha]{\overline{y}} \gamma H \underset{u}{\text{---}} \text{Ab}(u, s)$ where H is a free group of rank 2 and the rank 1 free extension of a centralizer $H *_u \text{Ab}(u, s)$ is generated by $\alpha, \gamma, \overline{x}$. Again γ and α cannot be conjugate, but u may be conjugate to one of them.

(F) The JSJ $F_{R(S)}$ has cyclic collapse $v \bullet \text{---} u$ and up to rational equivalence we can get the relative presentation

$$F_{R(S)} \approx F^\beta_\gamma \xrightarrow[\overline{y}]{\overline{x}} \delta_\epsilon H$$

where $\overline{x}$ and $\overline{y}$ are sent to stable letters and H is a free group of rank 2 generated by α, δ, ϵ .

In all cases the cyclic edge groups can be taken to be maximal cyclic in the vertex groups.

We finally note that this description in terms of relative presentations also gives a description of what *irreducible systems of equations* over F look like since these systems of equations are the relations of the groups. We make some remarks that could be seen as corollaries of this theorem.

Remark 1. If the JSJ of $F_{R(S)}$ has three vertices or three edges then one of the vertex groups is F and the other vertex groups are free of rank 2 or free abelian of rank 2.

Remark 2. If the JSJ of $F_{R(S)}$ has at least two nonabelian vertex groups, then one of them is free of rank 2.

Remark 3. If $\text{Rank}(F) = N$ and $b_1(F_{R(S)}) = N + 1$ then using Proposition 3.15 and looking at abelianized relative presentations we see that either $F_{R(S)}$ is a rank 1 free extension of a centralizer of F or $F_{R(S)}$ doesn't have any noncyclic abelian subgroups.

Direct inspection shows us that:

Corollary 2.7. *The height of the canonical analysis lattice relative to F (see Section 4 of [Sel01]) for $F_{R(S)}$ is at most 3, i.e. the lattice terminates at level L^2 .*

2.1 A description of Solutions

In [Ozh83], Ozhigov gives a description of the solution set of a system of equations in two variables over F . For each system of equations, his algorithm produces a finite collection of *forms*. Theorem 2.9 below gives us exactly the same description.

Definition 2.8. Let F be a free group. An expression of the form

$$f_1 p_1^{n_1} f_2 \cdots f_n p_n^{n_2} f_{n_1}$$

where the f_i and p_i lie in F and the n_i are variables in $\mathbb{Z}$ is called a *one level parametric word in F* . Inductively we define an *n -level parametric word* to be an expression of the form

$$f_1 p_1^{n_1} f_2 \cdots f_n p_n^{n_2} f_{n_1}$$

where $f_i \in F$, n_i are variables and the p_i are m -level parametric with $m < n$. Formally, a *parametric word* defines a subset of F .

Denote by $\Phi = \text{Stab}_{\text{Aut}(F(x,y))}([x, y])$. We have $\phi(x) = W(x, y)$, $\phi(y) = V(x, y)$. For a pair $(u, v) \in F \times F$ and $\phi \in \Phi$ as before we denote

$$\phi_x(u) = W(u, v), \phi_y(v) = V(u, v)$$

Theorem 2.9. *Let $S(x, y) = 1$ be any system of equations over F . Then the solutions of $S(x, y)$ in $F \times F$ are described by a finite number of families of pairs of the following form:*

- (a) (f, g) where f, g take arbitrary values in F .
- (b) $(P(F, r), Q(F, r))$ where P, Q are parametric words in $F * \langle r \rangle$ and r takes arbitrary values in F .
- (c) $(P(F), Q(F))$ where P, Q are parametric words in F
- (d) $(P(F, \phi_x(u), \phi_y(v)), Q(F, \phi_x(u), \phi_y(v)))$ where $u, v \in F$ are fixed and P, Q are fixed parametric words in $F \cup \{\phi_x(u), \phi_y(v)\}^{\pm 1}$, where ϕ takes arbitrary values in Φ .

Proof. Apply Theorem 1.26. Each family that is given corresponds to the solutions that factor through a branch of the Hom diagram. We also know what all the possible canonical automorphisms look like, in particular and non-surface automorphism will affect the variables in the parametric words.

From Theorem 2.6. it is easy to see the what $\text{Hom}_F(F_{R(S)}, F)$ is when $F_{R(S)}$ is freely decomposable, in particular all such Hom diagrams have one level. In particular they fall into category (a) and (b).

There is only one type fully residually F quotient of $F[x, y]$ that contains a QH subgroup, moreover by the description of Hom diagrams for such fully residually F groups given in [Tou07] (also deducible from [Ol'89, GK92, Lys88]) we see that any branch in the Hom diagram that contains a group with a QH subgroup will give a family of solutions describable as (d). $\square$

2.2 Examples and Questions

We give some examples of the groups given in Theorem 2.6. We first note that all the freely decomposable groups are easy to construct as quotients of $F[x, y]$. Examples of Theorem 2.6 (C) are easy to construct by taking extensions of centralizers of F or $F * \langle r \rangle$ or by taking a chain of extension of centralizers of height 2. The next few examples are more delicate.

Example 2.10 (Example of (D).I.1 of Theorem 2.6). Let $F = F(a, b)$. It is proved in [Tou07] that the group

$$\begin{aligned} G &= \langle F, x, y | [a^{-1}ba[b, a][x, y]^2x, a] = 1 \rangle \\ &\approx \langle F, x, y, t | [x, y]^2x = [a, b]a^{-1}b^{-1}at; [t, a] = 1 \rangle \end{aligned}$$

is freely indecomposable modulo F , that

$$G \approx F_1 *_{\langle u=w(x, y) \rangle} \langle x, y \rangle$$

where F_1 is a rank 1-extension of a centralizer of F . Moreover G is shown to be fully residually F by the F -embedding into the chain of extensions of centralizers

$$F_2 = \langle F, t, s \mid [t, a] = 1, [s, u] = 1 \rangle$$

where $u = [a, b]a^{-1}b^{-1}at$ via the mapping, $x \mapsto s^{-1}(b^{-1}t)s$ and $y \mapsto s^{-1}(b^{-1}ab)s$

Example 2.11 (Example of (E).I.2 of Theorem 2.6). We modify Example 2.10. Let $F = F(a, b)$ and let

$$F_1 = \langle F, s, t, r \mid [t, a] = 1, [s, b^{-1}ab] = 1, [u, r] = 1 \rangle$$

where $u = [a, b]a^{-1}b^{-1}at$. F_1 is a chain of extensions of centralizers. Let $x' = b^{-1}t, y' = b^{-1}ab$ and let $G = \langle F, r^{-1}x'r, sr \rangle$. Let $H = r^{-1}\langle x', y' \rangle r$ and consider $G \cap H$. We see that $(sr)^{-1}b^{-1}ab(sr) = r^{-1}b^{-1}abr$ so $H \leq G \cap H$, on the other hand letting $\bar{y} = (sr)$ and $\bar{x} = r^{-1}x'r$ and by Britton's lemma we have a splitting:

$$G = \tilde{F}^{b^{-1}ab} \xrightarrow[u]{\bar{y}} H$$

With $\tilde{F} = \langle F, t \rangle$ and H free of rank 2, not freely decomposable modulo edge groups.

Example 2.12 (Example of (D).III of Theorem 2.6). Let $F = F(a, b)$ and consider the chain of extensions of centralizers

$$F_2 = \langle F, s, t \mid [s, a] = 1, [t, (a^2(b^{-1}ab)^2)^s] = 1 \rangle$$

One can check that the subgroup $K \leq \langle F, s^{-1}bs, t \rangle$ has induced splitting:

$$F *_a \begin{array}{c} \curvearrowright^r \\ \gamma_H \gamma' \end{array} *_p Ab(p, t)$$

where $H = s^{-1}\langle a, b^{-1}ab \rangle s$, $\gamma = s^{-1}as$, $\gamma' = s^{-1}b^{-1}abs$, $r = s^{-1}bs$, and $p = (a^2(b^{-1}ab)^2)^s$. Moreover it is freely indecomposable, fully residually F and generated by two elements modulo F .

Compared to these, constructing an example of (F) of Theorem 2.6 is relatively easy.

2.2.1 Questions

Conspicuously absent from the list is an example of Theorem 2.6 (B). In fact the author has been unable to construct examples of such groups which leads me to the following conjecture:

Conjecture 2.13. There are no fully residually F groups generated by two elements modulo F such that their JSJ has only one vertex group, i.e. where the HNN extensions are *separated*.

Also absent from the examples are groups from Theorem 2.6 (C) such that the vertex group is not simply an extension of a centralizer or groups like in case (D).II.2 where u and q are *conjugate*. This motivates the following question:

Question 2.14. Consider the free group $F(a, b)$, is there a non primitive element $u \in F(a, b)$ and some element $w \in F(a, b)$ such that $\langle u, w \rangle \leq F(a, b)$ but for some conjugate pup^{-1} of u in $F(a, b)$ we have $\langle u, pup^{-1}, w \rangle = F(a, b)$

A positive answer would enable the construction of such groups, whereas a negative answer would probably exclude a few possibilities.

3 Tools for fully residually F groups:

Theorem 3.1. *Let $F_{R(S)}$ be fully residually F , then in particular it has the following properties.*

- *it is torsion free,*
- *it satisfies the CSA property, which implies commutation transitivity and implies that in any splitting, maximal abelian subgroups which do not split as HNN extensions must be elliptic,*
- *elements $g, h \in F_{R(S)}$ either commute or freely generate a free group of rank 2.*

Definition 3.2. A subgroup $K \leq G$ is said to have property CC (conjugacy closed) if for $k, k' \in K$

$$\exists g \in G \text{ such that } k^g = k' \Rightarrow \exists \tilde{k} \in K \text{ such that } k^{\tilde{k}} = k'$$

Lemma 3.3. $F \leq F_{R(S)}$ has property CC

Proof. Let $f, f' \in F$ we have a retraction $r : F_{R(S)} \rightarrow F$, then

$$f^g = f' = r(f') = r(f^g) = f^{r(g)}$$

□

We give some first tools for analyzing fully residually F groups.

3.1 Resolutions

For this section we consider the JSJ of $F_{R(S)}$. An extremely useful tool for studying the vertex groups and getting structural information is resolutions.

Definition 3.4. An epimorphism of $\rho : F_{R(S)} \rightarrow F_{R(S')}$ of fully residually F groups is called *strict* if it satisfies the following conditions on the cyclic JSJ splitting modulo F .

- For each abelian vertex group A , ρ is injective on the subgroup $A_1 \leq A$ generated by the boundary subgroups in A .
- ρ is injective on edge groups.
- The image of QH subgroups is nonabelian.
- For every rigid subgroup R , ρ is injective on the *envelope* $\tilde{R}$ of R , defined by first replacing each abelian vertex group with its boundary subgroups and letting $\tilde{R}$ be the subgroup of the resulting group generated by R and by the centralizers of incident edge groups.

Definition 3.5. A resolution of $F_{R(S)}$

$$\mathcal{R} : F_{R(S)} \xrightarrow{\pi_1} \cdots \xrightarrow{\pi_p} F_{R(S_p)} \xrightarrow{\pi_{p+1}} F$$

is a sequence of proper epimorphisms of fully residually F groups.¹ A *strict* resolution is a resolution such that all the epimorphisms are strict.

¹In [KM98b] this is called a *fundamental sequence*

The next result follow immediately from Theorem 1.26.

Theorem 3.6. [Sel01, KM98b] *If $F_{R(S)}$ is fully residually F then it admits a strict resolution.*

This next result, however, requires more work:

Lemma 3.7. [Sel01, KM98b] *The subset $\Phi_{\mathcal{R}} \leq \text{hom}_F(F_{R(S)}, F)$ of F -morphisms that factor through a strict resolution $\mathcal{R}$ F -discriminate $F_{R(S)}$.*

This next definition illustrates what is going on with non-free rigid vertex groups and why they eventually split along the strict resolution.

Definition 3.8. Let $R_1 \leq F_{R(S)}$ be a non free rigid subgroup and let $\beta \in R_1$ generate a boundary subgroup. We say that β *obstructs* R_1 if there is a splitting of R_1 in which β is hyperbolic. We say that an element of an abelian vertex group $A \leq F_{R(S)}$ is *exposed* if it does not lie in a boundary subgroup.

3.2 Some notions of complexity

We present two measures of complexity. The first one gives us bounds on the complexity of underlying graphs of groups:

Definition 3.9. Let $\mathcal{G}(X)$ be a graph of groups modulo F . We define the complexity $B(\mathcal{G}(X))$ to be following

$$B(\mathcal{G}(X)) = \text{Rank}(\pi_1(X)) + \sum_{v \in VX} \chi(v)$$

where

- $\chi(v) = 0$ if $F \leq G_v$.
- $\chi(v) = (\text{Rank}(G_v) - \sum \text{Rank}(B_i))$ if G_v is abelian and the B_i are the boundary subgroups in G_v .
- $\chi(v) = \max\{0, 2 - k\}$ if G_v is rigid and k is the number of incident edge groups.
- $\chi(v) = 2g$ if G_v is a QH subgroup and where g is the genus of the underlying surface.

Lemma 3.10. *Let H be a fully residually free group of rank two or greater, and let $\gamma \in H$ then the “one relator” quotient $H/(\gamma)$ is non trivial.*

Proof. If H is abelian, the result is clear. If H is nonabelian then H admits an epimorphism ψ onto a free group K of rank at least two, we have the following commutative diagram:

$$\begin{array}{ccc} H & \xrightarrow{\psi} & K \\ \downarrow & & \downarrow \\ H/(\gamma) & \xrightarrow{\bar{\psi}} & K/(\psi(\gamma)) = \bar{K} \end{array} \quad (3)$$

and the one relator group $\bar{K}$ is seen to be nontrivial by abelianizing. $\square$

Proposition 3.11. *The complexity $B(\mathcal{G}(X))$ gives a lower bound for the rank of $\pi_1(\mathcal{G}(X))$. Specifically, if $F_{R(S)} = \pi_1(\mathcal{G}(X))$ then $B(\mathcal{G}(X))$ cannot exceed the number of variables in S .*

Proof. Consider a quotient $\overline{G}$ of $G = \pi_1(\mathcal{G}(X))$ obtained by killing $\tilde{F}$, the vertex group containing F , and killing all the edge groups. $\overline{G}$ is therefore a free product, and Lemma 3.10 implies that the $\chi(v)$ indeed gives a lower bound for the rank of the image of G_v in $\overline{G}$. The lower bound for $\text{Rank}(G)$ now follows from Grushko's Theorem.

If G is a quotient of $F[x_1, \dots, x_n]$ then $\overline{G}$ is generated by $\langle x_1, \dots, x_n \rangle$, this proves the second part of the claim. $\square$

Corollary 3.12. *If $S = S(x, y)$ and $F_{R(S)} = \pi_1(\mathcal{G}(X))$ then $B(\mathcal{G}(X)) \leq 2$.*

Our second measure of complexity is the *first Betti number*.

Definition 3.13. The *first Betti number* of a group G denoted $b_1(G)$ is the rank of the torsion-free direct summand of the abelianization of G .

Since fully residually free groups are finitely presented it is easy to compute b_1 directly from the presentation. The fundamental group of a graph of groups $G = \pi_1(\mathcal{G}(X))$ with cyclic edge groups we have the following lower bound: for $T \subset X$ a maximal spanning tree we have

$$b_1(G) \geq \sum_{v \in T} b_1(G_v) - E \quad (4)$$

where E is the number of edges in T . If there is an epimorphism $G \rightarrow H$ then $b_1(G) \geq b_1(H)$. The following useful fact is obvious from a relative presentation:

Lemma 3.14. *Let $H < G$ be a rank n extension of a centralizer of H , then $b_1(G) = b_1(H) + n$.*

Proposition 3.15. *Let F be a free group of rank N and let G be a fully residually F F -group. Then $b_1(G) = N$ if and only if $G = F$.*

Proof. Suppose towards a contradiction that $G \neq F$ but $b_1(G) = N$. Then, by being fully residually F , there is a retraction $G \rightarrow F$. It follows that $b_1(G) \geq N$. By Corollary 1.24, G has D , a nontrivial JSJ decomposition. Let $F \leq \tilde{F} \leq G$ be the vertex group containing F , obviously $\tilde{F}$ is also fully residually F . By formula (4), $b_1(G) \geq b_1(\tilde{F})$ and if D has more than one vertex group then the inequality is proper which forces $b_1(G) > N$ – contradiction. We must therefore have that D is a bouquet of circles with a single vertex group $\tilde{F}$. By Lemma 3.3 if $\tilde{F} = F$ then the stable letters of the splitting D in fact extend centralizers of elements of F , so by Lemma 3.14 we have that $b_1(G) > b_1(F)$ – contradiction.

It follows that we cannot have $\tilde{F} = F$. We therefore look JSJ of $\tilde{F}$. Again we find that it must have a unique vertex group $\tilde{F}^1$. Combining Theorem 1.23, finite presentability of G , and the hierarchical accessibility result in [DP01] over the class of cyclic splittings gives a finite sequence of inclusions

$$\tilde{F} > \tilde{F}^1 > \dots \tilde{F}^r > \tilde{F}^{r+1} = F$$

where $\tilde{F}^{i+1}$ is the unique vertex group of the JSJ of $\tilde{F}^i$. Now, by assumption, we must have $N = b_1(F) \geq b_1(\tilde{F}^1) \dots \geq b_1(\tilde{F}^{r+1}) = N$ but we must have that $\tilde{F}^r$ has a splitting D^r that is a bouquet of circles with vertex group F , we saw that in this case we must have $b_1(\tilde{F}^r) > b_1(F)$ –contradiction. $\square$

3.3 Rational Equivalence

Definition 3.16. A map $f : F[x, y] \rightarrow F[x, y]$ induced by the mapping

$$\begin{aligned} f &\mapsto f; \text{ if } f \in F \\ x &\mapsto X(F, x, y) \\ y &\mapsto Y(F, x, y) \end{aligned}$$

is called a polynomial map.

Proposition 3.17. Let f be a polynomial map such that $f(x) = X(F, x, y)$ and $f(y) = Y(F, x, y)$ and let $S_1(F, x, y)$ be a system of equation in two variables. Let $S_2(F, x, y) = S_1(F, f(x), f(y))$. Then the map $\tilde{f} : F \times F \rightarrow F \times F$ given by

$$(a, b) \mapsto (X(F, a, b), Y(F, a, b))$$

restricts to a mapping $\tilde{f} : V(S_2) \rightarrow V(S_1)$ called a morphism of varieties.

Definition 3.18. If there are polynomial maps $f, g : F_{R(S)} \rightarrow F_{R(S)}$ such that $\tilde{f} \circ \tilde{g} = 1_{V(S_1)}$ and $\tilde{g} \circ \tilde{f} = 1_{V(S_2)}$ then we say that the systems of equations S_1 is *rationally equivalent* to S_2 , moreover we say the varieties $V(S_1)$ and $V(S_2)$ are *isomorphic*.

Theorem 3.19 ([BMR99] Corollary 9). $F_{R(S)}$ and $F_{R(S_1)}$ are F -isomorphic if and only if S and S_1 are rationally equivalent.

Corollary 3.20. Suppose that $F_{R(S)}$ can be expressed in generators $F, \bar{x}', \bar{y}'$ with relations $R(F, \bar{x}', \bar{y}')$, then the systems of equations $S(F, x, y)$ and $R(F, x, y)$ are rationally equivalent.

Proof. In the other generators we have $\bar{x} = X(F, \bar{x}', \bar{y}'), \bar{y} = Y(F, \bar{x}', \bar{y}')$. and we see that the F -map induced by $\bar{x} \mapsto X(F, \bar{x}', \bar{y}'), \bar{y} \mapsto Y(F, \bar{x}', \bar{y}')$ is an isomorphism:

$$\langle F, \bar{x}, \bar{y} \mid S \rangle \rightarrow \langle F, \bar{x}', \bar{y}' \mid R \rangle$$

The result now follows from Theorem 3.19. $\square$

4 Bass-Serre Theory Techniques

We first establish some notation; we will denote the commutator $x^{-1}y^{-1}xy = [x, y]$. For conjugation we will use the following convention:

$$\begin{aligned} x^w &= w^{-1}xw \\ {}^w x &= wxw^{-1} \end{aligned}$$

We use this convention since ${}^x({}^y w) = {}^{xy}w$. Our group will always act on a tree T from the left i.e. for all $g, h \in F_{R(S)}$ and for all $v \in T$ we have

$$ghv = g(hv)$$

it follows that if for any point $v \in T$, and for any $g \in F_{R(S)}$ we will have $\text{stab}(gv) = g(\text{stab}(v))g^{-1} = {}^g\text{stab}(v)$.

We will write $w = w(F, X)$, for example, to denote a *word*; i.e. an explicit product of symbols from F and $X^{\pm 1}$.

Definition 4.1. Let G act on a simplicial tree T without edge inversions and let $K \leq G$ act on T via the normal restriction. We will say that two vertices v, w of T are K -equivalent or of the same K -type, denoted $tp_K(v) = tp_K(w)$, if there is a $k \in K$ such that $kv = w$. We similarly define have K -types for edges of $T(K)$.

Theorem 4.2 (The Fundamental Theorem of Bass-Serre Theory). *If G acts on a simplicial tree T without edge inversions and the quotient $A = G \backslash T$ is a finite graph, then G splits as a the fundamental group of a graph of groups with underlying graph A . Vertex and edge groups correspond to stabilizers of vertices and edges of T (respectively) via the action of G .*

Conversely, if G splits as a graph of groups, then it acts on a simplicial tree T without inversions called the Bass-Serre tree.

Definition 4.3. An action of G on a tree T is said to be k -acylindrical if the largest subset of T fixed by an element of G has diameter k . A *splitting* of G is said to be k -acylindrical if the action of G on the induced Bass-Serre tree is k -acylindrical.

We note that if we take a splitting of $F_{R(S)}$ whose edge groups are maximal abelian in the vertex groups, then the splitting will be 1-acylindrical.

4.1 Induced splittings and $\mathcal{G}(A)$ -graphs

Suppose that G has a splitting D as the fundamental group of a graph of groups and let H be a subgroup of G . Then G acts on a tree T and H acts on the minimal H -invariant subtree $T(H) \subset T$ in such a way that H also splits as a graph of groups. We call this splitting D_H the *induced splitting* of H .

We now present the folding machinery developed in [KWM05], which is a more combinatorial version of the Stallings-Bestvina-Feighn-Dunwoody folding techniques in Bass-Serre theory, used to compute induced splittings. Although this machinery is technically demanding, it gives an alternative to normal forms when dealing with fundamental groups of graphs of groups which greatly simplifies the arguments of Sections 5 and 7.

4.1.1 Basic definitions

We follow [KWM05].

Definition 4.4. A $\mathcal{G}(A)$ -graph $\mathcal{B}$ consists of an underlying graph B with the following data:

- A graph morphism $[\cdot] : B \rightarrow A$
- For each $u \in VB$ there is a group B_u with $B_u \leq A_{[u]}$, called a vertex group.
- To each edge $e \in EB$ there are two associated elements $e_i \in A_{[i(f)]}$ and $e_t \in A_{[t(f)]}$ such that $(e^{-1})_i = (e_t)^{-1}$ for all $e \in EB$.

Convention 4.5. We will usually denote $G(A)$ -graphs by $\mathcal{B}$ and will assume that the underlying graph of $\mathcal{B}$ is some graph B .

When drawing these graphs we give vertices v the label $(B_v, [v])$. We give an edge e the label $(e_i, [e], e_t)$. We will say that an edge e is of *type* $[e]$.

Definition 4.6. Let $\mathcal{B}$ be a $\mathcal{G}(A)$ -graph and suppose that $e_1^{\epsilon_1}, \dots, e_n^{\epsilon_n}$; where $e_j \in EB, \epsilon_j \in \{\pm 1\}$; is an edge path of B . A sequence of the form

$$b_0, e_1^{\epsilon_1}, b_1, e_2^{\epsilon_2}, \dots, e_n^{\epsilon_n}, b_n$$

where $b_j \in B_{t(e_j^{\epsilon_j})}$ is called a $\mathcal{B}$ -path. To each b path we associate a label

$$\mu(p) = a_0 [e_1]^{\epsilon_1} a_1 [e_2]^{\epsilon_2} \dots [e_n]^{\epsilon_n} a_n$$


where $a_0 = b_0(e_1^{\epsilon_1})_i, a_j = (e_j^{\epsilon_j})_t b_j (e_{j+1}^{\epsilon_{j+1}})_i$ and $a_n = (e_n^{\epsilon_n})_t b_n$ which is a $\mathcal{G}(A)$ -path.

Definition 4.7. Let $\mathcal{B}$ be a $\mathcal{G}(A)$ -graph with a basepoint u . Then we define the subgroup $\pi_1(\mathcal{B}, u) \leq \pi_1(\mathcal{G}(A), [u])$ to be the subgroup generated by the $\mu(p)$ where p is a $\mathcal{B}$ -loop based at u .

Example 4.8. Let $G = \pi_1(\mathcal{G}(X), u) = A *_C B$. The underlying graph is

$$u \bullet \xrightarrow[e]{} \bullet v$$

with $X_u = A, X_v = B$ and $X_e = C$. Let $g = a_1 b_2 a_3 b_4 a_5$ where $a_i \in A$ and $b_j \in B$. Then the $\mathcal{G}(X)$ -graph



whose vertex groups are all trivial is such that $\pi_1(\mathcal{B}, u) = \langle g \rangle$

This example motivates a definition.

Definition 4.9. Let $g = b_0, e_1^{\epsilon_1}, b_1, e_2^{\epsilon_2}, \dots, e_n^{\epsilon_n}, b_n$ be some element of $\pi_1(\mathcal{G}(X), u)$. Then we call the based $\mathcal{G}(X)$ -graph $\mathcal{L}(g; u)$ a *g-loop* if $\mathcal{L}(g; u)$ consists of a cycle starting at u whose edges are all coherently oriented and have labels

$$(b_0, e_1^{\epsilon_1}, 1), (b_1, e_2^{\epsilon_2}, 1), \dots, (b_{n-2}, e_{n-1}^{\epsilon_{n-1}}, 1), (b_{n-1}, e_n^{\epsilon_n}, b_n)$$

respectively.

Definition 4.10. $(\mathcal{G}(A), v_0)$ be a graph of groups decomposition of $F_{R(S)}$. Let the $\overline{x}, \overline{y}$ -wedge, $\mathcal{W}(F, \overline{x}, \overline{y}; u)$, be the based $\mathcal{G}(A)$ -graph formed from a vertex v with label (F, v_0) and two attaching the loops $\mathcal{L}(\overline{x}; v_0)$ and $\mathcal{L}(\overline{y}; v_0)$.

It is clear that $\pi_1(\mathcal{W}(F, \overline{x}, \overline{y}; u), u) = \langle F, \overline{x}, \overline{y} \rangle = F_{R(S)}$. Usually we will omit the mention of the vertex u .

4.1.2 Moves on $\mathcal{G}(A)$ -graphs

Let $\mathcal{B}$ be a $\mathcal{G}(A)$ graph, with underlying graph B . We now briefly define the moves on $\mathcal{B}$ given in [KWM05] that we will use, we will sometimes replace an edge e by e^{-1} to shorten the descriptions:

- A0: **Conjugation at v :** For some vertex v , assume w.l.o.g. that for each edge incident to v , $i(e) = v$. For some $g \in A_{[v]}$ do the following: replace B_v by gB_vg^{-1} , for each edge such that $i(e) = v$ replace e_i by ge_i , and for each edge such that $t(e) = v$ replace e_t by e_tg^{-1} .
- A1: **Bass-Serre Move at e :** For some edge e , replace its label $(a, [e], b)$ by $(ai_e(c), [e], t_e(c^{-1})b)$ for some c in $A_{[e]}$.
- A2: **Simple adjustment at u on e :** For some vertex u and some edge e such that w.l.o.g $i(e) = v$, we replace the label $(a, [e], b)$ by $(ga, [e], b)$ where $g \in B_u$.
- F1: **Simple fold of e_1 and e_2 at the vertex u :** For a vertex u and edges e_1, e_2 such that w.l.o.g. $i(e_1) = i(e_2) = u$ but $t(e_1) = v_1 \neq t(e_2) = v_2$ but $[v_1] = [v_2]$, if e_1 and e_2 have the same label, then identify the edges e_1 and e_2 . The resulting edge has the same label as e_1 and the vertex resulting from the identification of v_1 and v_2 has label $\langle B_{v_1}, B_{v_2} \rangle$.
- F4: **Double edge fold (or collapse) of e_1 and e_2 at the vertex u :** For edges e_1, e_2 such that w.l.o.g. $i(e_1) = i(e_2) = u$, $t(e_1) = t(e_2) = v$, and $[e_1] = [e_2] = f$ if they have labels (a, f, b_1) and (a, f, b_2) respectively, and moreover $[e_1] = [e_2] = f$. Then we can identify the edges e_1 and e_2 , the image of these edges has label (a, f, b_1) and the the group B_v is replaced by $\langle B_v, b_1^{-1}b_2 \rangle$. We will also call such a fold a collapse from u towards v .

The moves F2 and F3 in [KWM05] are analogous to F1 and F4, respectively only they involve simple loops. However, because these moves only show up in the proof of Lemma 8.1, we do not describe them explicitly. We also introduce four new moves:

- T1: **Transmission from u to v through e :** For an edge e such that $i(e) = u$ and $t(e) = v$ with label $(a, [e], b)$. Let $g \in A_{[e]}$ be such that $ai_{[e]}(g)a^{-1} = c \in B_u$, then replace B_v by $\langle B_v, b^{-1}t_{[e]}(g)b \rangle$.
- L1: **Long range adjustment:** Perform a sequence of transmissions through edges $e_1, \dots, e_n$ followed by a simple adjustment that leaves unchanged the labels of the edges $e_1, \dots, e_n$. Finally replace all the modified vertex groups by what they were before the sequence of transmissions.

N1: **Nielsen move:** Replace a wedge $\mathcal{W}(F, \overline{x}, \overline{y})$ by $\mathcal{W}(F, \overline{x}', \overline{y}')$, where the triple $(F, \overline{x}', \overline{y}')$ is obtained from $(F, \overline{x}, \overline{y})$ via a Nielsen move modulo F .

S1: **Shaving move:** Suppose that u is a vertex of valence 1 such that $u = t(e)$ and $v = i(e)$, e has label $(a, [e], b)$ and $B_u = b^{-1}(t_{[e]}(C))b$, where $C \leq G_{[e]}$. Then delete the vertex u and the edge e and replace B_v by $\langle B_v, a(i_{[e]}(C))a^{-1} \rangle$.

Convention 4.11. Although formally applying a move to a $\mathcal{G}(A)$ -graph $\mathcal{B}$ gives a new graph $\mathcal{B}'$ unless noted otherwise we will denote this new $\mathcal{G}(A)$ -graph as $\mathcal{B}$ as well.

We regard the transmission above as the group B_u sending the element c to B_v through the edge e . When edge groups are *slender* in the sense of [DS99], multiple transmissions can be used instead of the edge equalizing moves F5-F6 in [KWM05]. N1 type moves are products of A0-A3, L1 and F1 type moves. We also notice that the moves T1, L1 S1 and N1 do not change the group $\pi_1(\mathcal{B}, u)$. Note that vertices of valence 1 with trivial group can always be shaved off.

4.1.3 The Folding Process

Definition 4.12. A $\mathcal{G}(A)$ -graph such that it is impossible to apply any moves other than A0-A2 is called *folded*.

This next important result is essentially a combination of Proposition 4.3, Lemma 4.16 and Proposition 4.15 of [KWM05].

Theorem 4.13. [KWM05] *Applying the moves A0-A2, F1-F4, and T1 to $\mathcal{B}$ do not change $H = \pi_1(\mathcal{B}, u)$, moreover if $\mathcal{B}$ is folded, then the associated data (see Definition 4.4) gives the groups decomposition of H induced by $H \leq \pi_1(\mathcal{G}(A))$*

This theorem implies the existence of a folding process. Consider the three following moves:

- **Adjustment:** Apply a sequence of moves A0-A2, L1, S1, and N1.
- **Folding:** Apply moves of type F1 and F4.
- **Transmission:** Apply a move of type T1.

First note that each folding decreases the number of edges in the graph, and that adjustments (except for shavings) are essentially reversible. In the folding process there is therefore a finite number of foldings and between foldings there are adjustments and transmissions. Transmissions enlarge the vertex groups and so in a sense increase the complexity of the $\mathcal{G}(A)$ -graph.

The goal of the next section is to find conditions that enable us to perform a maximal number of foldings without having to resort to transmissions. In a sense it will give us a folding process where each step decreases the complexity of the $\mathcal{G}(A)$ -graph, making it more tractable.

4.1.4 Transmission graphs and 1-acylindrical splittings

Suppose that $\mathcal{G}(A)$ is 1-acylindrical and the edge groups are abelian. Let $\mathcal{B}$ be a $\mathcal{G}(A)$ -graph.

Definition 4.14. We define the *pretransmission graph* B_0 of $\mathcal{B}$ to be the underlying graph B of $\mathcal{B}$ equipped with the following colouring:

- A vertex v such that B_v is nonabelian is *green*.
- A vertex v such that B_v is abelian is *yellow*.
- Otherwise edges and vertices are black.

Suppose that we performed a sequence of transmission $t_1, \dots, t_n$ on a $\mathcal{G}(A)$ graph $\mathcal{B}$. We will produce a analogous sequence of coloured graphs $B_1, \dots, B_n$. As follows for B_{i+1} start with B_i and make the following changes:

1. If for a vertex v , B_v is abelian and nontrivial, then colour it *yellow*.
2. If for a vertex v , B_v is non-abelian, then colour it *green*.
3. If the transmission move t_{i+1} is through the edge e , then colour e *red*.

Lemma 4.15. *Let $\mathcal{G}(A)$ be 1-acylindrical and let $\mathcal{B}$ be a $\mathcal{G}(A)$ -graph and let B_n be a transmission graph. Then after performing the corresponding sequence of transmission moves we have that for each black vertex u , B_u is trivial, for each yellow vertex v , such that w.l.o.g. $v = t(e)$, and the incoming edge e is labeled $(a, [e], b)$ we have $B_u \leq b^{-1}(t_{[e]}(G_{[e]}))b$.*

Proof. This follows immediately from 1-acylindricity of $\mathcal{G}(A)$ and the construction. $\square$

The usefulness of the next lemmas is that they enables us to control the vertex groups during the folding process.

Definition 4.16. A path in a transmission graph that consists of two red edges and a yellow vertex u is called a *cancellable path centered at u* .

Lemma 4.17. *Let $\mathcal{G}(A)$ be 1-acylindrical and let $\mathcal{B}$ be a $\mathcal{G}(A)$ -graph and suppose that after a sequence of transmissions $t_1, \dots, t_n$ the transmission graph B_n has a cancellable path centered at u . Then it is possible to perform a folding (F1 or F4) at u in $\mathcal{B}$ using only a Bass-Serre A1 move, and maybe a conjugation A0 move.*

Proof. If u is yellow, but has two (or more) adjacent red edges, then w.l.o.g. we have $i(e) = i(e') = u$ and that e and e' have labels $(a, [e], b)$ and $(a', [e], b')$ respectively. Let A denote the image of $G_{[e]}$ in $A_{[e]}$. The fact that both edges e' and e are red in particular imply that $aAa^{-1} \cap a'A(a')^{-1} \neq \{1\}$ which implies that $(a')^{-1}(aAa^{-1})a' \neq \{1\}$. Now 1-acylindricity implies malnormality of the edge groups, it therefore follows that

$$a'^{-1}a = \alpha \in A \Rightarrow a = a'\alpha$$

This means that in $\mathcal{B}$ all we need is to do a Bass-Serre move A1 so that the label of e' becomes $(a'\alpha, [e], \alpha^{-1}b')$ and then (for move F1) a conjugation move A0 so that e' has label $(a'\alpha, [e], b) = (a, [e], b)$ same as e . $\square$

Lemma 4.18. *Let $\mathcal{G}(A)$ be 1-acylindrical and let $\mathcal{B}$ be a $\mathcal{G}(A)$ -graph. Suppose that after a series of transmissions and adjustments it is possible to perform either move F1 or F4 at a yellow or black vertex u , of the corresponding transmission graph, then it is also possible to perform a folding move F1 or F4 at v after only a sequence of adjustments.*

Proof. Let $\mathcal{B}'$ be the result of applying all the transmission moves to $\mathcal{B}$. Suppose that it was impossible to perform the move F1 or F4 at u in $\mathcal{B}$ but that it is possible to it in $\mathcal{B}'$. Then it follows that $B'_u > B_u$ and that there is an A2 move that can now be performed at u so that F1 or F4 is now possible.

Suppose first that u is black then $B_u = B'_u = \{1\}$, which in particular implies that the move F1 or F4 could already have been applied to u in $\mathcal{B}$.

Suppose now that u is yellow and that in $\mathcal{B}'$ we can perform an A2 move changing the label of e , where $i(e) = u$ and then after performing a moves A1, A0 at $t(e)$ we can do either move F1 or F4 identifying e and e' . In particular e and e' had labels $(a, [e], b)$ and $(a', [e], b')$ respectively and after doing move A2 the labels became $(a', [e], b)$ and $(a', [e], b')$ respectively. We wish to show that instead we can change the label of e in the same way by performing a long range adjustment in $\mathcal{B}$ or moves A0, A1.

The only obstruction to long range adjustments is if both edges e and e' are either directed or red. By the construction of the transmission graph, a yellow vertex cannot have two incoming directed edges or an incoming edge and a red edge because that implies that the vertex is green. It follows that the only obstruction is if both edges e, e' are red, in which case Lemma 4.17 applies. $\square$

Definition 4.19. The transmission graphs obtained after performing all possible transmissions is called a *terminal transmission graph*.

4.2 When $F_{R(S)}$ splits as a graph of groups with two cycles

We suppose our group $F_{R(S)} = \langle F, \overline{x}, \overline{y} \rangle$ can be collapsed to a cyclic graph of groups modulo F with one vertex and two edges, i.e.

$$F_{R(S)} = {}_t \circlearrowleft \widetilde{F} \circlearrowright {}_s \quad (5)$$

where $F \leq \widetilde{F}$. We denote by s and t the corresponding stable letters.

It should be noted that the three stable letter case cannot occur, since killing the vertex group $F_{R(S)}$ yields a quotient that is a free group of rank 3 that is generated by two elements –contradiction.

Definition 4.20. For a generating set X and a word $W = W(X)$ in X , for a letter $x \in X$ we denote the exponent sum of x in W by $\sigma_x(W)$.

Definition 4.21. For a stable letter t from a splitting of G and a word an element $g \in G$. Let $\langle X|R \rangle$ be a presentation that represents the splitting, i.e. $t \in X$ and we have the appropriate relation. Let $g = W(X)$ where $W(X)$ is in normal form. We define the *exponent sum* $\sigma_t(g)$ of t in g as

$$\sigma_t(g) = \sigma_t(W(X))$$

It follows from Britton's Lemma that this quantity doesn't depend the choice of word chosen to represent g .

Definition 4.22. For $g \in G$, we define the (t, s) -signature $\text{sgn}_{(t,s)}(g)$ of g to be the pair of integers.

$$\text{sgn}_{(t,s)}(g) = (\sigma_t(g), \sigma_s(g))$$

It is customary to drop the subscripts.

We must be able to write $t = T(F, \bar{x}, \bar{y})$ and $s = S(F, \bar{x}, \bar{y})$ as words in our generators. On the other hand, by Britton's Lemma, for any word $V = V(F, \bar{x}, \bar{y})$ we have for $u = t, s$

$$\sigma_u(V) = \sigma_{\bar{x}}(V)\sigma_u(\bar{x}) + \sigma_{\bar{y}}(V)\sigma_u(\bar{y}) \quad (6)$$

Which means that we should be able to express vectors $(1, 0)$ and $(0, 1)$ as linear combinations of $\text{sgn}(\bar{x})$ and $\text{sgn}(\bar{y})$ in $\mathbb{Z}^2$, *in particular they are linearly independent as vectors.*

Lemma 4.23. Suppose $F_{R(S)}$ split as (5), then if some word $W(F, \bar{x}, \bar{y})$ lies in $\tilde{F}$ then it must have exponent sum 0 in both $\bar{x}$ and $\bar{y}$.

Proof. If $w = W(F, \bar{x}, \bar{y}) \in \tilde{F}$, then $\text{sgn}(w) = (0, 0)$. The result now follows immediately from equation (6) and the fact that $\text{sgn}(\bar{x})$ and $\text{sgn}(\bar{y})$ are linearly independent. $\square$

Lemma 4.24. Suppose $A \leq F_{R(S)}$ is a noncyclic abelian subgroup of $F_{R(S)}$. Then A cannot be generated by words $W_i(F, \bar{x}, \bar{y})$ such that for each W_i the exponent sums in $\bar{x}$ and $\bar{y}$ are zero.

Proof. Let $\mathcal{R}$ be a strict resolution with fully residually F groups $\{F_{R(S_i)}\}_{i \in I}$. Then for some $j \in I$ we must have that a summand $\langle r \rangle \leq A$ is *exposed* in the JSJ of $F_{R(S_j)}$. Without loss of generality $\pi(A) = A' \oplus \langle r \rangle$, where π is the composition of maps $\pi : F_{R(S)} \rightarrow F_{R(S_j)}$. We can now view $F_{R(S_j)}$ as an HNN extension

$$F_{R(S_j)} = \langle G, r \mid \forall a \in A', [r, a] = 1 \rangle$$

On one hand $\bar{x}$ and $\bar{y}$ are sent to elements with normal forms $\bar{x}'(H, r), \bar{y}'(H, r)$ on the other hand, r is in the image of A and by hypothesis we can write $r = R(F, \bar{x}'(H, r), \bar{y}'(H, r))$ where R has exponent sum zero in $\bar{x}'(H, r)$ and $\bar{y}'(H, r)$, but by Britton's Lemma $R(F, \bar{x}'(H, r), \bar{y}'(H, r)) = r$ must have exponent sum zero in r which is a contradiction. $\square$

Corollary 4.25. If $F_{R(S)}$ has a cyclic splitting modulo F with two cycles, then none of the vertex groups can contain noncyclic abelian subgroups.

4.3 The Weidmann Nielsen Technique for Groups acting on Trees

We present some of the techniques developed by Richard Weidmann in [Wei02]. Let G be a group and let T be a simplicial tree on which G acts.

Definition 4.26. Let $M \subset G$ be partitioned as

$$M = S_1 \sqcup \dots \sqcup S_p \sqcup H$$

where $H = \{h_1, \dots, h_s\}$. We say that M has *markings* $(S_1, \dots, S_p; H)$. We now have elementary Nielsen transformations on *marked sets*:

$T1$: Replace some S_i by S_i^g where $g \in M - S_i$.

$T2$: Replace some element $h \in H$ by $g_1 h g_2$ where $g_1, g_2 \in M - \{h\}$

Let

$$T_i = \{x \in T \mid \exists g \in \langle S_i \rangle \text{ such that } gx = x\} \cup \text{the minimal } \langle S_i \rangle\text{-invariant subtree}$$

An important subcase will be when the groups $\langle S_i \rangle$ have global fixed points. It is convenient to note that there is no difference if we replace an element a subset S_i by the subgroup $\langle S_i \rangle$. We can now formulate the main results in [Wei02].

Theorem 4.27. *Let M be a set with markings $(S_1, \dots, S_p; H)$. Then either*

$$\langle M \rangle = \langle S_1 \rangle * \dots * \langle S_p \rangle * F(H)$$

or by successively applying transformations $T1$ and $T2$ we can bring $(S_1, \dots, S_p; H)$ to a normalized marked set $\tilde{M} = \langle \tilde{S}_1, \dots, \tilde{S}_p, \tilde{H} \rangle$ such that one of the following must hold:

1. $\tilde{T}_i \cap \tilde{T}_j \neq \emptyset$, for some $i \neq j$.
2. $\exists h \in \tilde{H}, h\tilde{T}_i \cap \tilde{T}_i \neq \emptyset$
3. *There is some $h \in \tilde{H}$ that has a fixed point.*

This is especially useful to us if we let $F \leq S_1$. Notice that replacing transformation T_1 by T_1' given by:

- For S_1 , do nothing.
- If $g \notin S_i$ replace S_i by $S_i^{g^{-1}}$. If $g \in S_i$ do nothing.
- If $g \neq h$ replace h by $h^{g^{-1}}$. If $g = H$ do nothing.

We can arrange so that in Theorem 4.27 $\tilde{S}_1 = S_1$ and that the results still holds. We call these *Nielsen moves on marked sets modulo F* .

4.4 Tricks that work for fully residually F groups

Lemma 4.28. *Let $F_{R(S)}$ act on a simplicial tree T with maximal abelian edge stabilizers. Then $F \leq F_{R(S)}$ acts transitively on the edges of T_F (see Definition 4.26). In particular if we have that A is an edge stabilizer and that $F \cap A = \langle \alpha \rangle$ and there are $g \in F_{R(S)}, u \in F$ such that $u^g = \alpha^n$ then there is some $g' \in F$ such that $u^{g'} = \alpha^n$, this means that $g^{-1}g' \in A$*

Proof. Let e_1, e_2 be edges of T_F . Let $\text{stab}(e_1) \cap F = \langle \gamma \rangle$, $\text{stab}(e_2) \cap F = \langle \beta \rangle$. There is some $g \in F_{R(S)}$ such that $ge_1 = e_2$. It follows that

$$g\gamma g^{-1}e_2 = (g\gamma g^{-1})e_1 = e_2$$

which means that

$$g\gamma g^{-1} \in \text{stab}(e_2) \Rightarrow [g\gamma g^{-1}, \beta] = 1$$

Suppose first that $g\gamma g^{-1} \notin \langle \beta \rangle$, then $\langle \beta, g\gamma g^{-1} \rangle$ form a free abelian group of rank 2, but since $\gamma \in F$ we see that Lemma 4.24 applies and yields a contradiction. It therefore follows that $g\gamma g^{-1} = \beta^n$ for some $n \in \mathbb{Z} - \{0\}$. Since F has property CC it follows that there exists $g' \in F$ such that $g'\gamma g'^{-1} = \beta^n$, it follows that $(g')^{-1}g$ fixes e_1 . Since e_1 and e_2 were arbitrary, the result follows. $\square$

We now focus on the situation where the action of $F_{R(S)}$ on its Bass-Serre tree T is 1-acylindrical. Of particular interest to us is the situation in Theorem 4.27 when $T_F \cap \rho T_F \neq \emptyset$ and where $d(v_0, \rho v_0) \leq 2$ where v_0 is the vertex of T fixed by F . To simplify notation we will use relative presentations. The path from v_0 to ρv_0 contains either one or two edge types in which case the splitting of $F_{R(S)}$ contains the “subgraphs”

$$\tilde{F}^R \xrightarrow{t} {}^S H \quad \text{or} \quad \tilde{F} \text{---} C \text{---} H \quad (7)$$

with $F \leq \tilde{F}$.

Lemma 4.29. *Suppose that we have $F \leq \tilde{F} \leq F_{R(S)}$ and T as above, assume moreover that the action of $F_{R(S)}$ on T is 1-acylindrical. If there is some ρ such that $T_F \cap \rho T_F \neq \emptyset$, with $v_0 = \text{fix}(F)$, then there are $f_1, f_2 \in F$ such that either*

1. *the path $[v_0, \rho v_0]$ contains one edge G -type and $f_2 \rho f_1 = h \in H$; or*
2. *the path $[v_0, \rho v_0]$ contains two edge G -types and $f_2 \rho f_1 = th$ for some $h \in H$; which could be assumed to be the stable letter t if we change the presentation by conjugating a boundary monomorphism.*

Proof. We case 1. We first must have that T_F is not just a point. If necessary, we conjugate boundary monomorphisms so that the lift $\tilde{e}$ of the edge e in T is fixed by some element of F . Suppose we have that there is only edge G -type traversed on the path from v_0 to ρv_0 , as in Figure 1. Then we must have

$$\rho = a_2 b_1 a_1; a_i \in \tilde{F}, b_1 \in H$$

and looking at Figure 1 we see that there must be some $\kappa \in F$ that fixes the edge $a_2 \tilde{e}$. Let $C = \text{stab}_G(\tilde{e})$ be the subgroup stabilizer of the edge $\tilde{e}$. Then there is some $g \in F_{R(S)}$ such that ${}^g \kappa \in C$. By Lemma 4.28 we find that there is some $g' \in F$ such that ${}^g \kappa = {}^{g'} \kappa \in C \cap F$. Letting $g' = f_2 \in F$ we have ${}^{f_2} \kappa = \gamma \in \text{stab}(\tilde{e})$ so

$$\gamma^{f_2} \in \text{stab}(a_2 \tilde{e}) \Rightarrow f_2^{-1} \tilde{e} = a_2 \tilde{e} \Rightarrow f_2 a_2 \tilde{e} = \tilde{e}$$




Figure 1: The path from v_0 to ρv_0 only contains 1 edge G-type




Figure 2: The path from v_0 to ρv_0 contains 2 edge G-types

which means $f_2 a_2 = b' \in \text{stab}(te) \leq H$. This means that

$$f_2 \rho = f_2 a_2 b_1 a_1 = b' b_1 a_1; \quad b' b_1 \in H$$

so we may assume from now on that $a_2 = 1$.

We look again at Figure 1 and we see that the edge $b_1 \tilde{e}$ must lie in ρT_F and since $\rho = b_1 a_1$ we have:

$$\begin{aligned} & b_1 \text{stab}(e) \cap b_1 a_1 F \neq 1 \\ & \Rightarrow \text{stab}(e) \cap a_1 F \neq 1 \\ & \Rightarrow a_1^{-1} \text{stab}(e) \cap F \neq 1 \end{aligned}$$

Which is like saying that $\text{stab}(a_1^{-1} e) = a_1^{-1} C \cap F \neq 1$. Reusing the argument to get rid of a_2 we can find a $f_1 \in F$ such that $a_1 f_1 \in \text{stab}(\tilde{e}) \leq H$ which means that $\rho f_1 = b \in H$. This completes case 1 of the Lemma.

We now tackle case 2 of the Lemma. Consider Figure 2. Then we see that since we have $T_F \cap \rho T_F \neq \emptyset$ and that T_K has radius at most 1, we must have that $d(v_0, \rho v_0) \leq 2$ which means that

$$\rho = a_2 t b_1 a_1, \quad a_i \in \tilde{F}, b_1 \in H$$

and letting $R = \text{stab}_{\tilde{F}}(t), C = \text{stab}(\tilde{e})$ we can arrange, perhaps after conjugating boundary monomorphisms, that $F \cap C \neq 1 \neq F \cap R$. We can then find f_1, f_2 such that

$$f_2 \rho f_1 = f_2 a_2 t b_1 a_1 f_1 = t b'$$

as in case 1. The proof is almost exactly as in case 1. The only difference is that we get $f_2 a_2 = r \in R$ which means that we can “commute” it through the stable letter t to get the result. $\square$

5 Coordinate Groups that are freely decomposable modulo F

Proposition 5.1. *Suppose that $F_{R(S)} = \tilde{F} * H$, then*

$$\text{Rank}(\tilde{F}) + \text{Rank}(H) \leq N + 2$$

Proof. First note that the underlying graph of the splitting $F_{R(S)} = \tilde{F} * H$ consists of an edge and two distinct vertices. Let $\mathcal{G}(A)$ denote this graph of groups and let $\mathcal{B}$ be any $\mathcal{G}(A)$ -graph. Only the moves A0-A3, F1, and F4 can be applied.

Take $\mathcal{W}$ to be the wedge $\mathcal{W}(F, \overline{x}, \overline{y})$. Since $\pi_1(\mathcal{W}) = F_{R(S)}$ we have by Theorem 4.13 that $\mathcal{W}$ can be brought to a graph with a single edge and two distinct vertices. The underlying graph of $\mathcal{W}$ has two cycles and A has no cycles, which means that two collapses must occur. Moreover each collapse, maybe after applying F1 moves, either contributes a generator to H or to $\tilde{F}$. The result now follows. $\square$

Corollary 5.2. *If $F_{R(S)}$ is freely decomposable modulo F then either it is one generated modulo F or*

$$F_{R(S)} \approx \begin{cases} F * \langle x, y \rangle \\ F *_u \text{Ab}(u, t) * \langle x \rangle \\ F * \text{Ab}(x, y) \end{cases}$$

Proof. Apply Proposition 5.1 and Theorems 2.1 and 2.2. $\square$

6 Splittings with one nonabelian vertex group and at least one abelian vertex group

We prove item (C) of Theorem 2.6, i.e. we consider the case where the JSJ of $F_{R(S)}$ has abelian vertex groups but only one nonabelian vertex group $\tilde{F} \geq F$.

Lemma 6.1. *If the cyclic JSJ decomposition of $F_{R(S)}$ modulo F contains only one nonabelian vertex group then we need only consider the following*

possibilities

$$F_{R(S)} = \begin{cases} A_1 *_u \tilde{F} *_v A_2 \\ \tilde{F} *_u A \\ \textcircled{\tilde{F} \xrightarrow{u} A} \\ A_1^{\beta_1}_{\delta_1} \xrightarrow{\alpha_1} \tilde{F}_\gamma^\alpha \xrightarrow{\beta} A \\ t \left(\begin{array}{c} \mu \tilde{F}_\gamma^\alpha \xrightarrow{\beta} A \\ \tilde{F} \xrightarrow{\eta} A \end{array} \right) \\ \tilde{F} \xrightarrow{\quad} A \end{cases}$$

Where the vertex groups A, A_1, A_2 are abelian.

Proof. By the complexity bound given in Corollary 3.12, the only missing possibility is

$$F_{R(S)} = A_1 *_u \tilde{F}_\gamma^\alpha \xrightarrow{\beta} A$$

But by Corollary 3.12 and see that the $\text{Rank}(A_1) = 2$, which means that $A_1 *_u \tilde{F}$ can be seen as an extension of a centralizer, which is an HNN extension and therefore falls in one of the mentioned cases. $\square$

Proposition 6.2. *If the cyclic JSJ decomposition modulo F of $F_{R(S)}$ contains at least two vertex groups but only one of them, $\tilde{F}$, is non-abelian, then the only possibilities for the splitting are*

$$F_{R(S)} = A_1 *_u \tilde{F} *_v A_2, \tilde{F} *_u A, \text{ or } \textcircled{\tilde{F} \xrightarrow{u} A}$$

with A, A_1, A_2 abelian.

Proof. First suppose that the underlying graph of D , the cyclic JSJ decomposition modulo F , contains two cycles. Then we can collapse D to a double HNN extension of some group H , say with stable letters s, t . By Corollary 4.25, H cannot contain any non-cyclic abelian subgroups – contradiction. The underlying graph of D therefore contains at most one cycle.

The only case left to check is when $F_{R(S)} = \tilde{F}_\gamma^\alpha \xrightarrow{\beta} A$. Since $\tilde{F} \neq F$ we have that α, γ obstruct $\tilde{F}$. Let $\mathcal{R} : F_{R(S)} \xrightarrow{\pi_1} \cdots \xrightarrow{\pi_p} F_{R(S_p)}$

be a strict resolution. As long as β, α are unexposed (which means, $\tilde{F}$ can't split) we have that $\tilde{F}$ and A' are mapped monomorphically. On one hand the images of β and δ must be sent to powers of some common element in F since they commute. By Lemma 3.7 the associated $\Phi_{\mathcal{R}}$ -morphisms F -discriminate $F_{R(S)}$, β and δ must be sent to arbitrarily high powers via $\Phi_{\mathcal{R}}$ -morphisms. It follows that for some $F_{R(S_i)}$ in $\mathcal{R}$, say, β is exposed. At this point either $\tilde{F}$ splits or it doesn't, if it doesn't split $\tilde{F}$ lies in a nonabelian vertex group and α therefore fixes a vertex of nonabelian type, whereas β doesn't contradicting the fact that α and β are conjugate in $F_{R(S_i)}$. If $\tilde{F}$ splits in $F_{R(S_i)}$, we since γ and δ are conjugate, γ must be elliptic which means that α obstructed $\tilde{F}$ and therefore is hyperbolic in $F_{R(S_i)}$, but β is elliptic – contradiction. $\square$

Combining Proposition 6.2 and the following proposition proves most of item (C) of Theorem 2.6.

Proposition 6.3. *If $F_{R(S)} = \tilde{F} *_{\alpha} A_1$ or $F_{R(S)} = A_2 *_{\beta} \tilde{F} *_{\gamma} A_3$ and $\text{Rank}(A_1) \geq 3$ then $\tilde{F} = F$.*

Proof. This follows immediately from computing $b_1(F_{R(S)})$. $\square$

Case (C) of Theorem 2.6 is therefore proved.

7 Splittings with two or more nonabelian vertex groups

7.1 Maximal Abelian Collapses

Suppose that the JSJ decomposition of $F_{R(S)}$ contains at least two non-abelian vertex groups. The complexity bounds given in Section 3.2 do not give us bounds on the actual number of vertices in D . To have something more tractable to work with we take D and do the following (see Definition 1.11):

- (i) If any boundary subgroup $\langle \alpha \rangle = i_e(G_e)$ is a proper subgroup of a maximal abelian subgroup A , then do a folding move where we replace G_e by a copy of A .
- (ii) Ensuring that the resulting graph of groups always has at least two non abelian vertex groups perform sliding and collapsing moves until it is no longer possible to decrease the number of vertices or edges

In the end that the resulting graph of groups $\mathcal{G}(A)$ will have one of three possible forms:

$$\tilde{F} \text{ --- } H, \quad \tilde{F} \text{ --- } \text{---} H, \quad \text{or} \quad \tilde{F} \text{ --- } \text{---} H \quad (8)$$

where the vertex group $\tilde{F}$ contains F and boundary subgroups are maximal abelian in their vertex groups. Moreover, by item (ii) of our construction, distinct edges have distinct boundary subgroups. It follows that this splitting is 1-acylindrical. It also follows that only the moves given in Section 4.1.2 are applicable to $\mathcal{G}(A)$ -graphs.

Definition 7.1. We will call such a splitting a *maximal abelian collapse* of $F_{R(S)}$.

7.1.1 The strategy

To analyze the possibilities for $F_{R(S)}$, for each possibility given in (8) we will do the following:

1. Get the abelian collapse of the JSJ of $F_{R(S)}$, and use this splitting as the underlying graph of groups.
2. Start with the wedge $\mathcal{W} = \mathcal{W}(F, \overline{x}, \overline{y})$. Then using Theorem 4.27 and Lemma 4.29 we will simplify $\mathcal{W}$ by N1 moves (see Section 4.1.2) so that the loop $\mathcal{L}(\overline{x})$ is somehow simple.

3. We will then apply moves F1, F4, and L1 to simplify the graph as much as possible. It will turn out that the resulting $\mathcal{G}(A)$ -graph $\mathcal{B}$ will have the same underlying graph as $\mathcal{G}(A)$.
4. All that will remain to get a folded graph is to make some transmission moves, keeping track of these will tell us how the vertex groups are generated.
5. Finally, by arguing algebraically we will recover the original cyclic JSJ decomposition modulo F .

7.2 The one edge case

We consider the case where $F_{R(S)}$ splits as

$$F_{R(S)} = \tilde{F} *_A H \quad (9)$$

with $F \leq \tilde{F}$, A maximal abelian in both factors and H non abelian. Throughout this section $\tilde{F}, A, H$ will denote these groups.

Lemma 7.2. *Let $F_{R(S)}$ split as in (9). Using Nielsen moves on $(F, \bar{x}, \bar{y})$ modulo F we can arrange; conjugating boundary monomorphisms if necessary; so that $\bar{x}$ lies in either $\tilde{F}$ or H .*

Proof. Since we are assuming free indecomposability of $F_{R(S)}$ modulo F , we can apply Theorem 4.27. Let T be the Bass-Serre tree induced from the splitting (9). Let v_0 be the vertex fixed by $F \leq F_{R(S)}$. We start by looking at the marked generating set $(F; \{\bar{x}, \bar{y}\})$. We consider different cases.²

Case 1) T_F is a point: Since $F_{R(S)}$ isn't freely decomposable by Theorem 4.27 without loss of generality $\bar{x}$ must be elliptic. $T_{\bar{x}}$ is either a vertex or an edge

Case 1.1) $T_F \cap T_{\bar{x}} = \emptyset$: Consider the the marked generating set $(F, \langle \bar{x} \rangle; \{\bar{y}\})$ and apply Theorem 4.27 again. We now find that either, without loss of generality, $\bar{y}T_{\bar{x}} \cap T_{\bar{x}} \neq \emptyset$ or $\bar{y}$ is also elliptic.

The first case is impossible. Indeed, by 1-acylindricity $T_{\bar{x}}$ is either an edge or a point so for $\bar{y}T_{\bar{x}} \cap T_{\bar{x}} \neq \emptyset$ we must have that $\bar{x}$ fixes one of the endpoints of $T_{\bar{x}}$ which implies that $\bar{y}$ is elliptic.

If $\bar{y}$ is also elliptic then the trees $T_F, T_{\bar{x}}, T_{\bar{y}}$ cannot all be disjoint, otherwise $F_{R(S)}$ would be freely decomposable. If $T_{\bar{y}} \cap T_F \neq \emptyset$ then we can switch $\bar{x}$ and $\bar{y}$ and pass to Case 1.2. Otherwise $T_{\bar{y}} \cap T_{\bar{x}} \neq \emptyset$ then the tree $T_{\langle \bar{y}, \bar{x} \rangle}$ is either an edge or has radius 1. Passing to the marking $(F, \langle \bar{x}, \bar{y} \rangle; \emptyset)$ and applying Theorem 4.27 implies that $T_{\langle \bar{y}, \bar{x} \rangle}$ can be taken so that $T_F \cap T_{\langle \bar{y}, \bar{x} \rangle} \neq \emptyset$. Which means that both $\bar{x}$ and $\bar{y}$ can be brought into H .

Case 1.2) $T_F \cap T_{\bar{x}} \neq \emptyset$: Since we have the splitting (9) and by our assumptions on edge stabilizers we have that if $\bar{x}$ fixes v_0 then $\bar{x} \in \tilde{F}$ and we are done.

Case 2) T_F is not a point: Conjugating boundary monomorphism, we can arrange for some generator α of A to lie in F . We apply Theorem 4.27 and find that either $\bar{x}T_F \cap T_F \neq \emptyset$ or $\bar{x}$ is elliptic. In the former

²Our notation is such that e.g. Case 2.3.4 is a subcase of Case 2.3.

case, since T_F has radius 1 and $F \leq F_{R(S)}$ has property CC we can apply Lemma 4.29 to make $\bar{x} \in H$ and we are done. Otherwise $\bar{x}$ is elliptic and we consider the next case.

Case 2.1) $T_F \cap T_{\bar{x}} = \emptyset$: We consider the marked set $(F, \bar{x}; \{\bar{y}\})$ and we see that applying Theorem 4.27 we can either arrange for $\bar{y}$ to be elliptic or get that either $T_F \cap \bar{y}T_F \neq \emptyset$ or $T_{\bar{x}} \cap \bar{y}T_{\bar{x}} \neq \emptyset$. In Case 1.1 the latter possibility was seen to be impossible unless $\bar{y}$ was elliptic. If $\bar{y}T_F \cap T_F \neq \emptyset$, then we can apply Lemma 4.29 as for the previous case and obtain that $\bar{y} \in H$ and we are done.

The remaining possibility in this case is that $\bar{y}$ is elliptic and $T_{\bar{y}} \cap T_F = \emptyset$. For our group not to be freely decomposable we must have that $T_{\bar{y}} \cap T_{\bar{x}} \neq \emptyset$, moreover since both $\bar{x}, \bar{y}$ are elliptic the tree $T_{\langle \bar{x}, \bar{y} \rangle}$ must have radius 1. This is dealt with exactly as in the end of Case 1.1.

Case 2.2) $T_F \cap T_{\bar{x}} \neq \emptyset$: Then we have that, after perhaps changing the boundary monomorphisms that $\bar{x} \in H$ or $\bar{x} \in \tilde{F}$ and we are done. $\square$

Lemma 7.3. *Suppose $F_{R(S)}$ splits as (9) maximal abelian group A with $F \leq \tilde{F}$ and H non abelian. If $\bar{x}$ can be brought into the subgroup $\tilde{F}$ via Nielsen moves modulo F , then $F_{R(S)}$ is freely decomposable.*

Proof. Let $\mathcal{G}(X)$ be the graph of groups representing the splitting (9). We start with the wedge $\mathcal{W}(F, \bar{x}, \bar{y})$, and suppose that after a sequence of N1 moves, the element $\bar{x}$ lies in $\tilde{F}$. Then $F_{R(S)}$ can be represented with a $\mathcal{G}(X)$ -graph $(\mathcal{B}, v)$, which consists of a vertex v and a $\bar{y}$ -loop $\mathcal{L}(\bar{y}, v)$, where the vertex v has label $\langle F, \bar{x} \rangle$.

Since $\pi_1(\mathcal{B}, v) = F_{R(S)}$, by Theorem 4.13 we should be able to bring $\mathcal{B}$ to $\mathcal{G}(X)$. Now only the vertex group B_v of $\mathcal{B}$ is non trivial. We do our folding process, but only doing adjustments and foldings, in particular, doing moves only A0-A3, F1, F4, L1 and S1. If a collapse occurs then $\mathcal{B}$ doesn't have any cycles and B_0 has an extra yellow vertex. By doing S1 shaving moves we can assume that B_0 has at most one yellow vertex. By Lemma 4.18, as long as the terminal transmission graph has at most one green vertex or a cancellable path, we will always be able to perform one of the moves F1, F4, S1 after a some sequence of L1 and A0-A3 moves.

We see that each step of the way the terminal transmission graph of $\mathcal{B}$ has at most one green vertex or cancellable path, unless one of three possibilities, let v be the vertex such that $B_v \geq \langle F, \bar{x} \rangle$:

- (a) $\mathcal{B}$ has two vertices v, u and one edge e and after a transmission B_u is generated by at most two elements. The graph is then folded, but we see that $H = A * H'$ which implies free decomposability.
- (b) $\mathcal{B}$ has no cycles, three vertices and two edges. We assume that all shaving moves were performed. Then the only possibility for $\mathcal{B}$ is that it has endpoints v and u , and B_u is cyclic. But then if the transmission graph had any red edges then the vertex u could have been shaven off. It follows that it is impossible for the transmission graph to have two green vertices.
- (c) $\mathcal{B}$ consists of a cycle of length 2, with vertices v and u . This means that the vertex group B_u is trivial. For $\mathcal{B}$ to be folded, after a transmission we must be able to perform a F4 collapse move.

If the collapse is towards u then no transmissions are needed and B_u will be generated by some element and the edge group, which implies free decomposability. If the collapse is towards v then if the collapse can be done without doing transmissions first, we reduce to case (a).

Otherwise, before the collapse, do a conjugation move at u and two transmissions so that $B_u = \langle b_1 A' b_1^{-1}, A'' \rangle$ where $A', A'' \subset A$ and $b_1 \notin A$, we now do a simple adjustment and collapse, in the end we get that $B_u = H = \langle b_1 A' b_1^{-1}, A \rangle$. Which, by Lemma 7.16 implies free decomposability. $\square$


Lemma 7.4. *Suppose $F_{R(S)}$ splits as (9) and via Nielsen moves we were able to bring $\bar{x} \in H$. Then $F_{R(S)}$ is generated by $\langle F, \bar{x}, \bar{y}' \rangle$ where $\bar{y}'$ also lies in H .*

Proof. We start with the graph with one edge e and two vertices v, u with $B_v = F$ and $B_u = \langle \bar{x} \rangle$ and then at v attach the $\bar{y}$ -loop $\mathcal{L}(\bar{y}, v)$. Start our adjustment-folding process applying only moves A0-A3, F1, F4, L1, S1, as much as possible. By Lemma 4.18, as long as the transmission graph has at most two green vertices, we will always be able to perform one of the moves F1, F4, S1 after a some sequence of L1 and A0-A3 moves.

Suppose first that along the process avoiding transmissions, a collapse occurred, then $\mathcal{B}$ doesn't have any cycles, either $\mathcal{B}$ can be brought to a graph with one edge and two vertices in which case, applying Lemma 7.3 if $B_u = \langle \bar{x} \rangle$, the result follows.

Otherwise, after shaving, we may assume that B has three vertices u, v, w and two edges. We already have an edge e between u and v which means that the other edge f must be either between v and w ; in which case we can perform F1 at v without doing any transmissions; or f is between w and u ; in which case transmission from w to u implies that w can be shaven off, so we can do an F1 move at u without transmissions. Either way, by Lemma 7.3 we must have $B_v = F$ and $B_u = \langle \bar{x}, \bar{y}' \rangle$ and all that is needed to get a folded graph is transmissions, so the result holds.

Suppose now that no collapses occurred, but it is impossible to perform any moves of type F1 or F4 without doing transmissions. Then by Lemma 4.18, the transmission graph must have at least three green vertices, this can occur only if one of these cases occur:



We first tackle Case I. If after transmissions a move of type F1 is possible, then it is easy to see that such a move could have been done without transmission and this reduces to Case III. It therefore follows that the edges f, f' have to be collapsed. If the collapse is towards w , then no transmission moves are necessary, we get a graph with three vertices

with $B_w = \langle \overline{y}' \rangle$, $B_v = F$ and $B_u = \langle \overline{x} \rangle$, we note that if any transmissions from w or u to v are possible, then we can shave off these vertices and end up as in the case of Lemma 7.3. It therefore follows that we can perform a F1 folding move at v and replacing $\overline{y}'$ by a conjugate if necessary we get a graph with two vertices v, u with $B_v = F$ and $B_u = \langle \overline{x}, \overline{y}' \rangle$, all that remains to get a folded graph are transmission moves so the result follows.

Suppose now that the collapse was towards v then first we transmit towards w through both edges f and f' (if we only needed to transmit through one edge, we could have used a long range adjustment contradicting the assumption.) Now suppose $i(f) = i(f') = w$ and after a conjugation move we have that f and f' have labels $(c, [f], b)$ and $(1, [f], b')$ respectively. Moreover $c \notin A$, since otherwise we could have used a Bass-Serre move A1, and then collapsed. Then after the transmissions $B_w = \langle \alpha, c\alpha c^{-1} \rangle$ where $\langle \alpha \rangle = F \cap A$. To be able to make an F4 move, we must be able to make a simple adjustment which implies that $c \in \langle \alpha, c\alpha c^{-1} \rangle$. Let $c\alpha c^{-1} = \beta$, we have that $\langle \alpha, \beta \rangle$ freely generate a free group of rank 2. But if $c = W(\alpha, \beta)$, then we would have the relation

$$W(\alpha, \beta)\alpha W(\alpha, \beta)^{-1} = \beta$$

which is impossible.

We now consider Case II, as before it may reduce to Case III. Let g and g' be the edges between v and w . Suppose first that the collapse is towards u , but there had to be transmissions to w first, then; as in the previous paragraph; we can derive a contradiction. It follows that the collapse must be towards w , but in this case long range adjustments can be used instead of transmissions and simple adjustments and we revert to the no cycle case.

Finally, we consider Case III, since there can be no transmissions from u to v , we must have that the collapse is from v to u . Note that since F has property CC and that it is possible to transmit through both edges towards v , instead of doing so one can make a simple adjustment A2 at F and collapse towards u and the result holds. This means that only one edge can transmit, so it follows that the F4 move can simply be preceded by long range adjustments and A0, A2 moves. But then we have that $B_v = \langle \overline{x} \rangle$ and the only moves left are transmissions which imply that $H = \langle \overline{x} \rangle * A$ which implies free decomposability of $F_{R(S)}$. $\square$

From the previous lemmas we get.

Proposition 7.5. *If $F_{R(S)}$ is freely indecomposable and splits as in (9) with $F \leq \tilde{F}$. Then $F_{R(S)}$ can be generated by F and two elements $\overline{x}, \overline{y}' \in H$. Therefore, up to rational equivalence, we can assume that $\overline{x}, \overline{y}$ are sent into H .*

This next proposition enables us to revert back to a cyclic splitting.

Proposition 7.6. *Suppose that $F_{R(S)}$ has a splitting as in (9) then $F_{R(S)}$ admits a cyclic splitting*

$$\tilde{F}' *_{\langle p \rangle} H'$$

where either:

1. $\tilde{F}' = F$ and H is generated by $\langle p, \bar{x}, \bar{y} \rangle$, $p \in F$ i.e. H is a three generated fully residually free group (see Theorem 2.1).
2. $\tilde{F} = \langle F, p \rangle$ and $H' = \langle \bar{x}, \bar{y} \rangle$ with $p \in H'$ i.e. H' is free of rank 2.

Proof. We first consider when the amalgamating maximal abelian subgroup A in (9) is cyclic. We write $A = \langle p \rangle$ and $\tilde{F}' = \tilde{F}$, $H' = H$, then we apply Proposition 7.5 and we get that $\bar{x}, \bar{y} \in H$. Looking at normal forms we have that $\tilde{F} = \langle F, p \rangle$ and $H = \langle p, \bar{x}, \bar{y} \rangle$ if $p \in F$ then $\tilde{F} = F$ and H is three generated fully residually free. If $p \notin F$ then we must have by normal forms that $p \in \langle \bar{x}, \bar{y} \rangle$ and it follows that $H = \langle \bar{x}, \bar{y} \rangle$.

We now consider the case where A in (9) is not cyclic. By Proposition 7.5 we can assume that $\bar{x}, \bar{y} \in H$ and it follows that $\tilde{F} = \langle F, A \rangle$. $\tilde{F}$ is fully residually F but not equal to F so $\tilde{F}$ has a nontrivial cyclic splitting modulo F and A must be elliptic. By Theorem 2.2, the only possibility for $\langle F, A \rangle$ is

$$\tilde{F} = F *_p A$$

which means that $F_{R(S)}$ admits the splitting:

$$F_{R(S)} = F *_p (A *_A H)$$

and as before we get that $H = \langle \bar{x}, \bar{y}, p \rangle$ is a three generated fully residually free group. $\square$

For the next result, we need:

Theorem 7.7 (Main Theorem of [Bau65]). *Let $w = w(x_1, x_2, \dots, x_n)$ be an element of a free group F freely generated by $x_1, x_2, \dots, x_n$ which is neither a proper power nor a primitive.³ If $g_2, g_2, \dots, g_n, g$ are elements of a free group connected by the relation*

$$w(g_1, g_2, \dots, g_n) = g^n \quad (n > 1)$$

then the rank of the group generated by $g_1, g_2, \dots, g_n, g$ is at most $n - 1$.

Lemma 7.8. *Suppose $F_{R(S)}$ splits as*

$$\tilde{F} *_u (\langle u \rangle \oplus \langle \gamma \rangle) *_\gamma \langle \bar{x}, \bar{y} \rangle$$

then $F_{R(S)}$ is freely decomposable, in particular, $\langle \gamma \rangle$ must be a free factor of $F(\bar{x}, \bar{y})$.

Proof. There are solutions $f : F_{R(S)} \rightarrow F$ such that the restriction of f to $F(\bar{x}, \bar{y})$ is a monomorphism since $F(\bar{x}, \bar{y})$ is a two generated nonabelian subgroup of a fully residually free group. We fix a solution f and denote by $H \leq F$ the image of $F(\bar{x}, \bar{y})$. H is free of rank 2 but we see that in the ambient group $\bar{\gamma} = f(\gamma)$ is a proper power, i.e. $\bar{\gamma} = u^n$ and n can be made arbitrarily large via Dehn twists. Theorem 7.7 applies and the assumption that $\gamma \in F(\bar{x}, \bar{y})$ is not a proper power (by malnormality of boundary subgroups) and not primitive (by free indecomposability modulo F) force the image of $F(\bar{x}, \bar{y}) \rightarrow F$ to be cyclic contradicting injectivity. $\square$

³An element in a free group is termed primitive if it can be included in a set of free generators

Proposition 7.9 (Theorem 2.6 (D) item 1). *If $F_{R(S)}$ has a QH subgroup then the only possibility is*

$$F_{R(S)} = F *_{u=p} \langle x, y, p | [x, y] = p \rangle$$

Proof. Note that by Proposition 3.11 and Corollary 3.12 there can be at most one MQH subgroup which must have genus at most 1, and the underlying graph of the graph of groups must be simply connected. The only suitable surface is the torus with one puncture. Again using the complexity bound in Corollary 3.12 the only possible cyclic JSJ splittings modulo F are of the form:

$$F_{R(S)} = \tilde{F}' *_{\underbrace{q H_1 *_{a_2} \dots *_{a_k} H_k}_{H}} *_{p} Q$$

where Q is the MQH subgroup and $F \leq \tilde{F}'$. Consider the splitting

$$(\tilde{F}' *_{q} H) *_{p} Q$$

p lies inside a maximal abelian subgroup A then we fold if necessary to get

$$(\tilde{F}' *_{q} H) *_{A} (A *_{p} Q)$$

we now apply Proposition 7.5 to get $\bar{x}, \bar{y} \in (A *_{p} Q)$. First suppose that A is noncyclic abelian, then we get $\tilde{F} = \tilde{F}' *_{q} H = \langle F, A \rangle$ which as in the proof of Proposition 7.6 must be $\tilde{F} = F *_{q} A$ so we get that $F_{R(S)} = F *_{q} A *_{p} Q$. If $p \neq q$ then, by Lemma 7.8, $F_{R(S)}$ is freely decomposable modulo F . If $p = q$ then if $\mathcal{G}(X)$ is the corresponding graph of groups with vertex groups F, A, Q then we find that $B(\mathcal{G}(X)) \geq 3$ which contradicts Corollary 3.12.

The possibility $A = \langle p \rangle$ remains. Proposition 7.5 gives us that $F_{R(S)}$ is generated by F and by $\bar{x}, \bar{y} \in Q$ and by Proposition 7.6 we have either $p \in F$ or $p \notin F$, in the former case the result follows i.e. we have that $Q = \langle \bar{x}, \bar{y}, p \rangle$ and $\tilde{F} = F$. In the latter case we have $\tilde{F} = \langle F, p \rangle$ and we must have $p \in \langle \bar{x}, \bar{y} \rangle$. By Theorem 2.2 and by free indecomposability modulo F we must have that $\langle F, p \rangle = \tilde{F} \approx \langle F, t | t^{-1} u t = u \rangle$. I.e $\tilde{F}$ contains a free abelian group of rank 2. On the other hand p has exponent sum 0 in both $\bar{x}$ and $\bar{y}$, so by Lemma 4.24 we have a contradiction. $\square$

Lemma 7.10 (Lemma 2.10 of [Tou07]). *Let G be a free group of rank 2 and let $w \in G$ be non primitive, and not a proper power. Then the only possible almost reduced (see Definition 1.18), hairless nontrivial cyclic splittings of G as the fundamental group of a graph of groups with w elliptic is as an HNN extensions*

$$G = \langle H, t | t^{-1} p t = q \rangle; p, q \in H - \{1\}$$

where $w \in H$ and H is another free group of rank 2. Moreover we have that $H = \langle p \rangle * \langle q \rangle$ which implies $G = \langle p, t \rangle$.

Proposition 7.11. *Suppose that $F_{R(S)}$ splits as*

$$F *_{\langle p \rangle} H \tag{10}$$

and $H = G *_q Ab(q, t)$ is a rank 1 free extension of a centralizer of a free group of rank 2. Then (10) refines to

$$F *__{\langle p \rangle} G *_q Ab(q, t) \quad (11)$$

Proof. This situation corresponds to item 1 of Proposition 7.6. We need to show that p is conjugable into G . Suppose towards a contradiction that this was impossible.

We first consider the possible cyclic JSJ splittings of $F_{R(S)}$, since q must be elliptic, any induced cyclic splitting of H must induce a cyclic splitting of G modulo q . G is a free group of rank two so, by Lemma 7.10, G may only split further as an HNN extension $G = \langle G', s \mid s^{-1}us = u' \rangle$ modulo q .

Now consider the cyclic JSJ splitting of H modulo p , by Lemma 7.8 p is not conjugate to a power of t . By our assumption and, if necessary, by replacing G with a vertex group G' containing p and q we have that H has no nontrivial cyclic splittings modulo p . On the other hand H is not free and there is an H -discriminating family of F -homomorphisms that send p to some fixed element of F , we must therefore have that H has a cyclic splitting modulo p —contradiction. $\square$

We note that this proof did not exclude the case where $p=q$.

Proposition 7.12. *Suppose that $F_{R(S)}$ splits as*

$$\tilde{F} *__{\langle p \rangle} H \text{ or}$$

where H is free of rank 2 and suppose moreover that H splits further as an HNN extension

$$H = \langle G, t \mid t^{-1}\mu t = \mu' \rangle$$

modulo p . Then we have that p and μ cannot be conjugate in H .

Proof. By Lemma 7.10, we must have that $H = \langle \mu, t \rangle$, if α and μ are conjugate in H then by conjugating boundary monomorphisms and doing Tietze transformations we see that $F_{R(S)} = \tilde{F} * \langle t \rangle$. $\square$

The same argument yields:

Proposition 7.13. *Suppose that $F_{R(S')}$ splits as*

$$F *__{\langle p \rangle} H *_u Ab(u, t)$$

with H free of rank 2 and suppose moreover that H splits further as an HNN extension

$$H = \langle G, t \mid t^{-1}\mu t = \mu' \rangle$$

modulo p and u , then we cannot have that both p and u are conjugate to μ

Corollary 7.14. *If $F_{R(S)}$ is freely indecomposable and the maximal abelian collapse of its cyclic JSJ decomposition modulo F is a free product with amalgamation. Then all the possibilities for the JSJ of $F_{R(S)}$ are given in item (D) of Theorem 2.6*

7.3 The two edge case

We now consider the case where, after “folding and sliding”, the splitting of $F_{R(S)}$ has underlying graph

$$X = v \bullet \begin{array}{c} \xrightarrow{e} \\ \xleftarrow{f} \end{array} \bullet u \quad (12)$$

and to which we give the relative presentation:

$$\tilde{F}^B \xrightleftharpoons[A]{t} {}^C H \quad (13)$$

Where $F \leq \tilde{F} = X_u, H = X_v$ and A, B, C are maximal abelian and conjugacy separated (in their vertex groups). In particular we have that the action of $F_{R(S)}$ on the corresponding Bass-Serre tree is 1-acylindrical. We now prove the lemmas that will enable us to prove item (E) of Theorem 2.6.

Lemma 7.15. *If $F_{R(S)}$ splits as in (13). Using Nielsen moves on $(F, \bar{x}, \bar{y})$ modulo F we can arrange; conjugating boundary monomorphisms if necessary; so that $\bar{x}$ either lies in $\tilde{F} \cup H$ or $\bar{x} = t$.*

Proof. We first observe that $F_{R(S)}$ cannot be generated by elliptic elements w.r.t. the splitting (13). We assume free indecomposability modulo F and apply Theorem 4.27 to the marked generating set $(F; \{\bar{x}, \bar{y}\})$. Let T be the Bass-Serre tree of the splitting. Let $v_0 \in T$ be the vertex fixed by $F \leq F_{R(S)}$.

Suppose that T_F is a point. Then by Theorem 4.27 we are forced to have that $\bar{x}$ can be brought to an elliptic element and that we can arrange $T_{\bar{x}} \cap T_F \neq \emptyset$ or $\bar{y}T_{\bar{x}} \cap T_F \neq \emptyset$ it follows that we can bring $\bar{x} \in \tilde{F}$. Suppose now that T_F is not a point either

1. $T_F \cap \bar{x}T_F \neq \emptyset$. Then we can apply Lemma 4.29 and get that $\bar{x}$ is either in H or $\bar{x} = th, h \in H$; or
2. $\bar{x}$ is elliptic. Then we use Theorem 4.27 on the marked set $(F, \bar{x}; \{\bar{y}\})$. $\bar{y}$ cannot also be elliptic.

If $T_F \cap T_{\bar{x}} \neq \emptyset$ then if $v_0 \in T_{\bar{x}}$ then we can assume that $\bar{x} \in \tilde{F}$. If $\bar{x}$ fixes a vertex w' adjacent to v_0 , then we can assume that $\bar{x} \in \text{stab}(w')$. By Lemma 4.28 we either have that $\bar{x}$ can be brought into H or tHt^{-1} , after possibly changing the spanning tree, the result follows.

If $T_F \cap \bar{y}T_F \neq \emptyset$ then as before we can arrange to that $\bar{y} = th$ and interchanging $\bar{x}$ and $\bar{y}$ the result will follow. The remaining case is $T_{\bar{x}} \cap \bar{y}T_{\bar{x}} \neq \emptyset$. We note, however, that $T_{\bar{x}}$ has at most one $F_{R(S)}$ -type of edge, but since $\bar{y}$ have exponent sum 1 in the stable letter, we find that the path ρ connecting $T_{\bar{x}}$ and $\bar{y}T_{\bar{x}}$ must contain both edge types. It follows that $T_{\bar{x}} \cap \bar{y}T_{\bar{x}} = \emptyset$. So we must have $T_F \cap \bar{y}T_{\bar{x}} \neq \emptyset$ which reduces to an earlier possibility.

□

Lemma 7.16. *Let $A, B \leq G$ be two abelian subgroups of a fully residually free group G such that for some $a \in A, b \in B$ we have $[a, b] \neq 1$. Then we have*

$$\langle A, B \rangle = A * B$$

Proof. Let $w = a_1 b_2 a_3 \dots b_n$ be product of non trivial factors $a_i \in A$ and $b_j \in B$ with perhaps the exception that a_1 or b_n are trivial. Since G is fully residually free there exists a map of G into F such that all the nontrivial a_i, b_j as well as some commutator $[a, b], a \in A, b \in B$ do not vanish. We have that the a_i are sent to powers of some element $u \in F$ and the b_j are sent to powers of some $v \in F$. It follows that the homomorphic image of w is sent to a freely reduced word in u and v , and since $u, v \in F$ do not commute they freely generate a free subgroup of F . It follows that w is not sent to a trivial element. $\square$

Lemma 7.17. *If the vertex group H in the splitting (13) is generated by conjugates of its boundary subgroups, i.e. $H = \langle A^{h_1}, C^{h_2} \rangle$, $h_i \in H$, then $F_{R(S)}$ is freely decomposable modulo F .*

Proof. Without loss of generality, by conjugating boundary subgroups we may assume, $h_1, h_2 = 1$. Take the presentation of $F_{R(S)}$ from the splitting, since by Lemma 7.16 $H = A * C$ we can use a Tietze transformation to get rid of C . The resulting group will have a relative presentation

$$\langle \tilde{F}, t | \emptyset \rangle = \tilde{F} * \langle t \rangle$$

$\square$

Lemma 7.18. *Suppose that $F_{R(S)}$ splits as in (13) and that $\bar{x} \in \tilde{F}$, then $F_{R(S)}$ is freely decomposable modulo F .*

Proof. It is clear that, perhaps after Nielsen moves modulo $\langle F, \bar{x} \rangle, \bar{y}$ can be sent to an element with exactly one occurrence of the stable letter t . A and C are not conjugate, which implies that $B \cap \langle F, \bar{x} \rangle = B' \neq 1$ and that the occurrences of t cancel in some product

$$\zeta^{-1} \beta \zeta$$

for some $\beta \in B'$. Let $C' = (B')^{\bar{y}}$, we have that C' is conjugate into H . Let $A' = \langle F, \bar{x} \rangle \cap A$.


To derive free decomposability it is enough to assume that $C' \leq H$. $C' \cap A = 1$ which means that $\langle C', A \rangle = C' * A$ and in particular that $\langle C, A' \rangle \cap A = A'$ which means that $H \cap \langle F, \bar{x}, \bar{y} \rangle = \langle A, C' \rangle$, which by Lemma 7.17 implies that $F_{R(S)}$ is freely decomposable. $\square$

Lemma 7.19. *Suppose that $\bar{x} \in H$, then we can arrange so that $F_{R(S)}$ is generated by F and elements $\bar{x} \in H$ and some $\bar{y}' = th, h \in H$.*

Proof. We first note that if $\bar{x}$ is conjugate into an edge group, then we can assume $\bar{x} \in \tilde{F}$, which leads to a contradiction. The hypotheses imply that $F_{R(S)}$ is the fundamental group of a $\mathcal{G}(X)$ -graph $\mathcal{B}$ obtained by taking an edge labelled, say $(1, e, 1)$, with endpoints v and u , attaching the $\bar{y}$ -loop $\mathcal{L}(\bar{y}, v)$, and setting $B_v = F, B_u = \langle \bar{x} \rangle$.


The pretransmission graph B_0 has the green vertex v and the yellow vertex u . We start our folding process using only moves A0-A3,F1,F4,L1,S1. Note that if a F4 collapse move occurs, then the underlying graph will be simply connected, which is impossible. By Lemma 4.18, this will be able to continue folding and adjusting as long as the terminal transmission graph has at most two green vertices or a cancellable path, which must be the case if, after shaving, there are more than four vertices.

Suppose that $\mathcal{B}$ has only four vertices, noting that this must fold to a graph like (12) using F1 moves we see (exchanging the labels e and f , if necessary) that the only possibilities after doing S1 moves are:




where here the edges are labelled by their type. If we look more closely at the possible transmissions, we see that the vertices u_1, v_1 will always be yellow in transmission graphs, so we can continue folding without using transmissions.

Suppose now that $\mathcal{B}$ has only three vertices, then the only possibilities after doing S1 moves are



Since the only folding that can occur is a F4 move at v or u , we see that we can use moves A0-A4,L1 and then make an F1 move.

It follows that $\mathcal{B}$ can be brought to a graph of the form:



with $B_v = F$ and $B_u = \langle \bar{x} \rangle$. Moreover our assumption that $\bar{x} \in H$ meant that $a_1 = b_1 = 1$. $F_{R(S)}$ is therefore generated by $F, \bar{x} \in H$ and $\bar{y}' = f, b_2, e, a_2$ with $a_2 \in \tilde{F}$ and $b_2 \in H$, by conjugating boundary monomorphisms we can assume that $\bar{y}' = t$. $\square$


Lemma 7.20. *Suppose that $\bar{x} = t$, then we can arrange so that $F_{R(S)}$ is generated by $F, \bar{x}$ and some $\bar{y}' \in H$.*

Proof. The hypotheses imply that $F_{R(S)}$ is the fundamental group of a $\mathcal{G}(X)$ -graph $\mathcal{B}$ obtained by taking two edges with labels $(1, e, 1)$ and $(1, f, 1)$ with common endpoints v and u , setting $B_v = F, B_u = \{1\}$, and attaching the $\bar{y}$ -loop $\mathcal{L}(\bar{y}, v)$.

The pretransmission graph B_0 has only the green vertex v , but the terminal transmission graph has green vertices u and v . We start our folding process using only moves A0-A3,F1,F4,L1,S1 as much as possible.


By Lemma 4.18, this will be able to continue as long as the terminal transmission graph has at most two green vertices or a cancellable path.

Suppose first that a F4 collapse occurred. Then $\mathcal{B}$ has one cycle. After shaving, the only possibilities (interchanging labels e and f , if necessary) so that the terminal transmission graph has more than two green vertices are



We note that in this case, one can always use moves A0-A3, and L1 so that F1 can be performed. In these cases we either get a graph with $B_v = \langle F, \overline{y'} \rangle, B_u = \{1\}$ or $B_v = F, B_u = \{\overline{y'}\}$. To get a folded graph, all that remains are transmissions, but note that in the former case, Lemma 7.17 will imply free decomposability modulo F , in the latter case the result follows.

Suppose that no collapses occurred, but the terminal transmission graph has more than two green vertices and no cancellable paths. Then the $\mathcal{B}$ has at most 4 vertices the only possibility with 4 vertices is something of the form:




Either an F1 type fold can be performed at u or v , in which case we need only make moves A0-A3, L1 before, or the fold occurs at v_1 or u_1 . In these latter cases (noting that all vertex groups except B_v are trivial) we see that if either u_1 or v_1 become green then before doing the F1 type fold, we will only need to use moves A0-A3. It follows that we can continue without transmissions.

Suppose now that $\mathcal{B}$ has three vertices then the possibilities are:



If the next fold will be of type F1, then it will occur at either v or u and moves A0-A3, L1, will be sufficient, and we get a graph with two vertices and three edges. If the next fold is a collapse, first note that, by the exact same arguments as in Lemma 7.4 Case I, if it is possible to get to collapse towards either v or u , then this must be possible without using transmissions first. Either collapse will reduce the situation to (14).

Suppose now that $\mathcal{B}$ has two distinct vertices and three edges, the possibilities are



where the remaining edge is marked only by its type. Note that these cases are symmetric. Suppose the third edge has label (a, e, b) with $a \in \bar{F}$ and $b \in H$. First note that if it is possible to transmit through both e -type edges from v to u , because F has property CC this means that there is some $a' \in F$ such that $a'a \in X_e$ which means that after an A1 Bass-Serre move we can make a F4 collapse towards u .

The remaining case is that one cannot transmit through both e -type edges from v to u , but then this means that we can make a long range adjustment to either bring (a, e, b) to $(a, e, 1)$ or bring $(1, e, 1)$ to $(1, e, b)$, and collapse.

If the collapse is towards v , then $B_v = \langle F, \bar{y}' \rangle$ and $B_u = \{1\}$, or $B_v = F$ and $B_u = \langle \bar{y}' \rangle$. All that remains to be done to get a folded graph is transmissions, in the latter case the result follows, in the former case we can derive free decomposability modulo F . $\square$

All these lemmas combine to give:

Proposition 7.21. *If we have the splitting described in (13) then we have that $F_{R(S)}$ is generated by the stable letter t and some element $\bar{y}' \in H$. Therefore, up to rational equivalence, we can assume that $\bar{x} = t$ and $\bar{y}$ is sent into H .*

The next proposition enables us to revert to a cyclic splitting.

Proposition 7.22. *Suppose that $F_{R(S)}$ has a splitting as in (13), then $F_{R(S)}$ admits one of the two possible cyclic splittings:*

1. $F_{R(S)} = F^\beta \xrightarrow[\alpha]{\bar{x}} H'$ where $H' = \langle \alpha, \bar{x}^{-1}\beta\bar{x}, \bar{y} \rangle$ is a three generated fully residually free group.
2. $F_{R(S)} = \tilde{F}^\beta \xrightarrow[\alpha]{\bar{x}} H'$ where $\tilde{F}$ is a rank 1 free extension of a centralizer of F and H' is generated by $\alpha, \bar{y}$. Moreover H' may not split further as an HNN extension.

Proof. Let $\tilde{F}, A, B, C, H, t$ be as in (13). By Proposition 7.21 we can assume that $\bar{x} = t$ and $\bar{y} \in H$. We can always assume that either $F \cap A \neq \{1\}$ (otherwise we could derive free decomposability modulo F .)

Suppose first that $F \cap A = \langle \alpha \rangle$ and $F \cap B = \{1\}$. To ensure free indecomposability modulo F we need there to be some $\gamma \in \langle \alpha, \bar{y} \rangle$ such that $\bar{x}\gamma\bar{x}^{-1} = \beta \in \tilde{F} \cap B$. Now by Theorem 2.2 if $\langle F, \beta \rangle \neq F * \langle \beta \rangle$ then we must have $\langle F, \beta \rangle = \langle F, t[u, t] = 1 \rangle$ for some $u \in F$. If u is not conjugate to α in F then $F_{R(S)}$ has a cyclic splitting as in item 2. If u and α are conjugate in F , then we can assume that $\alpha = u$, so then the group A in (13) is noncyclic abelian of rank 2. We study the maximal abelian subgroup $C \leq H$ we already had that $\gamma(\bar{y}, \alpha) \in C$ if $C \leq H$ is not cyclic then there must be some $\gamma_1 \in \langle \bar{y}, A \rangle$ such that γ and γ_1 do not lie in a common cyclic subgroup and which satisfies the relation

$$[\gamma, \gamma_1] = 1 \tag{15}$$

however by Lemma 7.16 we have

$$\langle A, \bar{y} \rangle = A * \langle \bar{y} \rangle \tag{16}$$

which means that (15) is impossible. It follows that $\tilde{F} = \langle F, \beta \rangle$. This gives the cyclic splitting:

$$F_{R(S)} = \langle F, t \rangle \xrightarrow[\alpha]{\overline{\alpha}} \gamma \langle \alpha, \overline{y} \rangle$$

Suppose now towards a contradiction that $\langle \alpha, \overline{y} \rangle$ split further as an HNN extension:

$$\langle \alpha, \overline{y} \rangle = \langle K | p^t = q \rangle; p, q \in K$$

then we have

$$F_{R(S)} = \tilde{F}_\alpha^\beta \xrightarrow[\alpha]{\beta'} K_q^p \rangle$$

Then we can collapse this splitting to a double HNN extension, and applying Lemmas 4.23 and 4.24 we see that $\tilde{F}$ cannot contain any noncyclic abelian subgroups –contradiction.

Suppose now that $F \cap A = \langle \alpha \rangle$ and $F \cap B = \langle \beta \rangle$: Then we first consider $H'' = \langle F, \alpha, \gamma \rangle$ where $\gamma = \overline{x}^{-1} \beta \overline{x}$. Suppose that A and B are non cyclic, then we see that

$$\langle F, A, B \rangle = A *_\alpha F *_\beta B$$

since $\langle F, A, B \rangle$ is fully residually F . which means that letting $H' = \langle H'', A, B \rangle$ we get the cyclic splitting

$$F_{R(S)} \approx F^\beta \xrightarrow[\alpha]{\overline{\alpha}} \gamma H'$$

and looking at normal forms and words in $\{F, \overline{x}, \overline{y}\}$ we see that $H' = \langle \alpha, \gamma, \overline{y} \rangle$. In particular we have a splitting as in item 1. *Having exhausted all the possibilities, the result holds* $\square$

Proposition 7.23. *Suppose that $F_{R(S)}$ splits as*

$$F^\beta \xrightarrow[\alpha]{\overline{\alpha}} \gamma H$$

and suppose moreover that H splits as an HNN extension

$$H = \langle G, s \mid s^{-1} \mu s = \mu' \rangle$$

modulo α, γ . Then α and γ cannot both be conjugate to μ in H .

Proof. After conjugating boundary monomorphisms and applying Lemma 7.10, it is easy to get Tietze transformations that exhibit a free decomposition. $\square$

Proposition 7.24. *Suppose that $F_{R(S)}$ splits as $F_{R(S)} = F^\beta \xrightarrow[\alpha]{\overline{\alpha}} \gamma H'$ where H' is a rank 1 free extension of a centralizer of a free group H of a free group of rank 2. Then this splitting can be refined to*

$$F_{R(S)} = F^\beta \xrightarrow[\alpha]{\overline{\alpha}} \gamma H \xrightarrow[u]{\quad} Ab(u, s)$$

such that α and γ are not conjugate and such that u is not conjugate to both α, γ . This cyclic splitting cannot be further refined.

Proof. Noting that the elements α and $\gamma = \overline{x}^{-1}\beta\overline{x}$ of H' are conjugate into F enables us to use the arguments of Proposition 7.11 to show that the elements α and γ must be conjugable into H' .

If u is conjugate to both α, γ , then it is clear that the group is freely decomposable modulo F .

Finally, if this splitting could be refined, then the only possibility by Lemma 7.10 is that H refines to an HNN extension, but then we could apply Lemma 4.24, contradicting the existence of a non-cyclic abelian subgroup. $\square$

All these Propositions imply:

Corollary 7.25. *If $F_{R(S)}$ is freely indecomposable and the maximal abelian collapse of its cyclic JSJ decomposition modulo F has two edges. Then all the possibilities for the JSJ of $F_{R(S)}$ are given in item (E) of Theorem 2.6*

7.4 The three edge case

We now consider the case where, after “folding, sliding, and collapsing”, the splitting of $F_{R(S)}$ has underlying graph

$$X = v \bullet \begin{array}{c} \xrightarrow{e} \\ \xrightarrow{f} \\ \xleftarrow{g} \end{array} \bullet u \quad (17)$$

and to which we give the relative presentation:

$$\tilde{F}_D^A \begin{array}{c} \xrightarrow{s} \\ \xrightarrow{C} \\ \xleftarrow{t} \end{array} \begin{array}{c} B \\ E \end{array} H \quad (18)$$

Where $F \leq \tilde{F} = X_v, H = X_u$ and A, B, C, D, E are maximal abelian in their vertex groups, hence the splitting is 1-acylindrical. Note that by Corollary 4.25, $F_{R(S)}$ cannot contain any noncyclic abelian subgroups, in particular the subgroups A, B, C, D, E must all be cyclic.


Lemma 7.26. *Let $F_{R(S)}$ split as in (18). Using Nielsen moves on $(F, \overline{x}, \overline{y})$ modulo F we can arrange; conjugating boundary monomorphisms if necessary so that $\overline{x} = t$.*

Proof. Since we are assuming free indecomposability of $F_{R(S)}$ modulo F , we can apply Theorem 4.27. Let T be the Bass-Serre tree corresponding to the splitting (17). We note that neither $\overline{x}$ nor $\overline{y}$ can be brought to elliptic elements w.r.t. the splitting (17). Let v_0 be the vertex fixed by F . W.l.o.g. we have $T_F \cap \overline{x}T_f \neq \emptyset$. 1-acylindricity implies that $d(v_0, \overline{x}v_0) = 2$. It follows w.l.o.g. that $\overline{x}$ is of the form a_1, f, b_1, e^{-1}, a_2 where $b_1 \in H, a_1, a_2 \in \tilde{F}$, by Lemma 4.29 we can arrange so that $\overline{x} = f, b, e^{-1}$, and conjugating boundary monomorphisms enables us to assume that $\overline{x} = f, e^{-1} = s^{-1}$ in terms of the relative presentation (18). $\square$

Lemma 7.27. *Suppose that $F_{R(S)}$ splits as in (18) and that $\overline{x} = t$, then $F_{R(S)}$ is generated by F, t , and s*


Proof. The hypotheses imply that $F_{R(S)}$ is the fundamental group of a $\mathcal{G}(X)$ -graph $\mathcal{B}$ obtained by taking two edges with labels $(1, e, 1)$ and $(1, f, 1)$ with common endpoints v and u , setting $B_v = F, B_u = \{1\}$, and attaching the $\overline{y}$ -loop $\mathcal{L}(\overline{y}, v)$.

Again we start our adjustment-folding process, using only moves A0-A2, L1, F1, and F4. Moreover we see that F4 collapses are forbidden since they reduce the number of cycles in the underlying graph. As long as there are strictly more than 4 vertices, the terminal transmission graph will only have two green vertices or a cancellable path. Suppose that the terminal transmission graph has only 4 vertices then, interchanging e and f if necessary, and noting that $\text{sgn}_s(\overline{y}) = 1$, the only possibilities are:




where the edges are marked by their type. In all three cases, though, we see that the vertices w_1, v_1 in the terminal transmission graph must be yellow, so we can continue our adjustment-folding process.

If there are only three vertices then we only the possibilities:



and the last fold is of type F1 at either u or v and in particular no transmissions are needed. We get that B is given by




Where the edges labelled e and f have labels $(1, e, 1)$ and $(1, f, 1)$ respectively. In the end we have that $F_{R(S)}$ is generated by F, t and some element $\overline{y}' = a_1, g, b_1, e^{-1}$ where $b_1 \in H$ and $a_1 \in \widetilde{F}$. After conjugating boundary monomorphisms, we may assume that $\overline{y}' = s$. $\square$

Corollary 7.28. *If $F_{R(S)}$ is freely indecomposable and the maximal abelian collapse of its cyclic JSJ decomposition modulo F has three edges. The JSJ of $F_{R(S)}$ is what is described in item (F) of Theorem 2.6*

Proof. All we need to show is that the vertex groups are F and a free group of rank 2.

By the two previous lemmas we have that $F_{R(S)}$ is the fundamental group of the $\mathcal{G}(X)$ -graph



with $B_v = F$ and $B_u = \{1\}$. To get a folded graph, all that are needed are transmissions. We also saw that the edge groups are cyclic. Suppose first that the only possible transmission is from u to v through e , then by 1-acylindricity, it is impossible for there to be any further transmissions from u back to v through the other edges.

Suppose that now there were transmissions possible only from v to u through edges e and f . So as not to have free decomposability modulo F , we must have a transmission from u to v through g . We note that the boundary subgroups associated to the edges e, f must be maximal cyclic because they lie in F , it then follows that there are no further possible transmissions and the graph is folded. In particular we find that $B_u = \tilde{F} = \langle F, \alpha \rangle$ where α is the element transmitted from H to $\tilde{F}$. $F_{R(S)}$ is freely indecomposable only if $\tilde{F} \neq F * \langle \alpha \rangle$, but by Theorem 2.2 the only other possibility for $\langle F, \alpha \rangle$ is $F *_u Ab(u, t)$, which is impossible since $F_{R(S)}$ has no noncyclic abelian subgroups.

It therefore follows that $\tilde{F} = F$ and H is a free group of rank 2 generated by its boundary subgroups. $\square$

8 Splittings with one vertex group

We now consider the situation where $F_{R(S)}$ has a cyclic JSJ decompositions modulo F , which yields relative presentations:

$$\begin{array}{c} \curvearrowright \\ \beta \tilde{F} \beta' \end{array} \quad (19)$$

$$t \left(\begin{array}{c} \alpha \\ \alpha' \tilde{F} \beta' \end{array} \right)^s \quad (20)$$

8.1 First Classifications

We give a first description of the vertex group. Note that if $\tilde{F} = F$ then the splittings (19) (20) must be extensions of centralizers of F .

Lemma 8.1. *Suppose $F_{R(S)}$ has the cyclic JSJ decomposition modulo F (20) then either $\tilde{F} = F$, or the following hold:*

1. *One of the vertex groups, say $\langle \alpha \rangle$, is conjugate into F .*
2. *$\tilde{F}$ is two generated modulo F .*
3. *The elements α and β are not conjugate in $F_{R(S)}$*
4. *$\tilde{F}$ has no abelian subgroups*
5. *The splitting is 1-acylindrical.*
6. *Up to rational equivalence we can assume $\overline{x} = t$ and $\overline{y} = t$*
7. *If $\tilde{F}$ is freely indecomposable, then its JSJ has two vertices and at most two edges. Moreover one of the vertex groups is F .*

Proof. The fact that $\tilde{F}$ cannot contain any abelian subgroups follows immediately from Corollary 4.25. We are assuming that $\tilde{F} \neq F$ which means that $\tilde{F}$ has a nontrivial cyclic $D_{\tilde{F}}$ splitting modulo F . It follows that α

and β must obstruct this splitting. On the other hand we have free indecomposability of $F_{R(S)}$, and it is impossible for both $\bar{x}$ and $\bar{y}$ to be elliptic in this splitting (linear algebra on exponent sums of stable letters.) It follows that Theorem 4.27 forces T_F to have edges, i.e. w.l.o.g. $\alpha \in F$.

Suppose now that α and β were conjugate. $\alpha \in F$ would imply that α', β, β' are elliptic in $D_{\bar{F}}$, contradicting the fact that α, β obstruct the splitting $\langle \delta \rangle_{\bar{F}}$. We have thus shown that the edge groups are maximal abelian and that α, β are not conjugate, it follows that the splitting (20) is 1-acylindrical.

We apply Theorem 4.27 the marked generating set $(F; \{\bar{x}, \bar{y}\})$. We can arrange via Nielsen moves so that $T_F \cap \bar{x}T_F \neq \emptyset$, by 1-acylindricity we have $d(v_0, \bar{x}v_0) < 2$ where $v_0 = \text{fix}(F)$. Since β is not conjugate into F we have that path from v_0 to $\bar{x}v_0$ has only t -type edges, and since $\text{sgn}_{(t,s)}(\bar{x}), \text{sgn}_{(t,s)}(\bar{y})$ are a basis of $\mathbb{Z}^2$ (see Definition 4.22) we have that $\bar{x} = f_1 t f_2, f_i \in \bar{F}$, so conjugating boundary monomorphisms we can assume that $\bar{x} = t$.

$F_{R(S)}$ is given by the $\mathcal{G}(X)$ -graph $\mathcal{B}$ with one vertex labelled v with label $(\langle \bar{F}, \alpha' \rangle, [v])$, an edge e with label $(1, [e], 1)$, i.e. $\mathcal{L}(\bar{x}, v)$; and the $\bar{y}$ -loop $\mathcal{L}(\bar{y}, v)$. By 1-acylindricity we see that we can bring $\mathcal{B}$ to a graph with one vertex and two edges without any transmissions. In order to get a folded graph, we must finally to transmit through the other edge so that the new label of v is $(\langle F, \alpha', \beta' \rangle, [v])$. It follows that $F_{R(S)}$ is generated by F, s, t and $\tilde{F} = \langle F, \alpha', \beta' \rangle$.

Finally, suppose that the JSJ of $F_{R(S)}$ had only one vertex. Then since $\tilde{F} \neq F$ the JSJ is nontrivial. Eventually the monomorphic image of $\tilde{F}$ will split in some term $F_{R(S')}$ of a strict resolution of $F_{R(S)}$. In particular, $\tilde{F}$ is generated by elliptic elements F and α' . Moreover since the induced splitting of $\tilde{F}$ is as an HNN extension, then $F_{R(S')}$ must also split as an HNN extension, say with stable letter z . Since F and α' are elliptic, $\beta \in \langle F, \alpha' \rangle$ must have exponent sum 0 in z . On the other hand since $\tilde{F} = \langle F, \alpha', \beta' \rangle$ splits as an HNN extension, β' is forced to have exponent sum 1 in z , which is impossible since β' and β are conjugate in $F_{R(S')}$.

Since $\tilde{F}$ cannot have any abelian subgroups, previous classifications imply that the JSJ of $F_{R(S)}$ has exactly two vertex groups one of which is F . Moreover in the $\tilde{F}$ is generated by F and α' which are elliptic and some β' which may or may not be elliptic. Now a simple exponent sum argument excludes the possibility of a JSJ with more than two edges. $\square$

Corollary 8.2. *Suppose that $F_{R(S)}$ has cyclic JSJ modulo F :*

$$\begin{array}{c} \alpha \\ \alpha' \end{array} \tilde{F} \xrightarrow{u} A$$

with A abelian, then α and u are not conjugate and moreover either α or u are conjugate into F .

Lemma 8.3. *If $F_{R(S)}$ splits as (19) and $\tilde{F} \neq F$ then $\beta, \beta' \notin F \leq \tilde{F}$ and $\tilde{F}$ is 2 generated modulo F , in particular $\tilde{F} = \langle F, \bar{x}, \beta' \rangle$. Moreover the splitting is 1-acylindrical and, up to rational equivalence, we can assume that $\bar{y} = t$.*

Proof. Suppose towards a contradiction that $\beta \in F$, then since $\tilde{F} \neq F$ so $\tilde{F}$ has a nontrivial cyclic splitting modulo F . It follows that β' must also be elliptic in $D_{\tilde{F}}$, in which case we can “refine” the splitting (19), contradicting the fact that it’s a cyclic JSJ modulo F . 1-acylindricity now follows from the fact that β, β' are hyperbolic in a cyclic splitting of $\tilde{F}$ modulo F , which means that β, β' cannot lie in noncyclic abelian subgroups.

We now let $F_{R(S)}$ act on the Bass-Serre tree corresponding to (19) and consider the marked generating set $(F; \{\bar{x}, \bar{y}\})$ we have that T_F must be a point, so by Theorem 4.27 $\bar{x}$ can be sent to an elliptic element and can be sent into $\tilde{F}$ via Nielsen moves modulo F . It follows that we must have $\beta \in \langle F, \bar{x} \rangle$. And since we must have $T_{\langle F, \bar{x} \rangle} \cap \bar{y}T_{\langle F, \bar{x} \rangle} \neq \emptyset$ by 1-acylindricity we easily conclude $\bar{y} = f_1 t^{\pm 1} f_2$ for $f_1, f_2 \in \tilde{F}$, conjugating boundary monomorphisms, we may therefore assume that $\bar{y} = s$ and we have $\tilde{F} = \langle F, \bar{x}, \beta' \rangle$. $\square$

8.2 The case where $\tilde{F}$ is not freely decomposable modulo F

In order to be able to say more about $\tilde{F}$ we pass to a quotient of $F_{R(S)}$. In a strict quotient $\pi : F_{R(S)} \rightarrow F_{R(S')}$, the restriction $\pi|_{\tilde{F}}$ is injective but the image $\pi(\tilde{F})$ will have a nontrivial induced cyclic splitting modulo F . We will identify $\tilde{F}$ with its image in $F_{R(S')}$. The author wishes to acknowledge that many ideas in this section, in particular Lemma 8.10 as well as its effect on the first Betti number come from the proof of Proposition 4.3 of [Sel01], i.e. the construction of the so-called cyclic analysis lattice.

Convention 8.4. We will first consider the case where the JSJ of $F_{R(S)}$ has one vertex and one edge. Hence from now until Section 8.2.3 $F_{R(S)}$ will be assumed to have a JSJ with one edge and one vertex.

Lemma 8.5. *Let $\pi : F_{R(S)} \rightarrow F_{R(S')}$ be a strict quotient. Then there are no cyclic splittings of $F_{R(S)}$ modulo $\tilde{F}$.*

Proof. Suppose towards a contradiction that was not the case. Then we can obtain a cyclic splitting of $F_{R(S')}$

$$A *_C B \text{ or } \begin{array}{c} \curvearrowright \\ C \\ A_{C'} \end{array}$$

such that $\tilde{F} \leq A$. π is surjective. If $F_{R(S')}$ split as an amalgam then we would have

$$\pi(\bar{y})^{-1} \beta \pi(\bar{y}) = \beta \quad (21)$$

where β and $\bar{y}$ lie in A , and $\pi(\bar{y})$ must not lie in A . If B is abelian of rank 3 then $A = F$ and we must have that $\tilde{F} = F$. If B is abelian of rank 2 then we will consider this splitting as an HNN extension and tackle it in the next case. We can therefore assume that B is nonabelian and by malnormality of the centralizer of C in both factors we see that (21) is impossible.

Suppose now that $F_{R(S')}$ split as an HNN extension then $\bar{y}$ must have exponent sum 1 in the stable letter r . In particular, by Britton's lemma we see that β and β' must be conjugate into C and C' , that $A = F_{R(S')} \cap \langle F, \pi(\bar{x}), \pi(\bar{y}) \rangle$ must be equal to $\tilde{F}$ and so that that conjugating boundary monomorphisms we can bring $\pi(\bar{y})$ to r , contradicting the fact that π is a proper surjection. $\square$

Corollary 8.6. *Any cyclic splitting of $F_{R(S')}$ induces a cyclic splitting of $\tilde{F}$.*

Corollary 8.7. *If $\tilde{F}$ is freely indecomposable modulo F , then so must be $F_{R(S')}$.*

Definition 8.8. Let e be an edge in the cyclic JSJ splitting of $\tilde{F}$ modulo F . We say that e is *visible* in a one edge splitting D of $F_{R(S')}$ if the edge group associated to e is conjugate into the edge group of D .

Definition 8.9. We will call the conjugacy classes of a boundary subgroup boundary an *edge class*.

8.2.1 Controlling induced splittings

The proofs in this section are very technical. Their goal is to obtain Corollary 8.14. Using this corollary we will be able to apply the following Lemma:

Lemma 8.10. *If $F_{R(S')}$ has a one edge cyclic D splitting and if $\tilde{F}$ has an induced one edge cyclic splitting $D_{\tilde{F}}$, then $F_{R(S')}$ can be obtained from $\tilde{F}$ by adding an element $\sqrt[n]{\eta}$: an n^{th} root of the generator η of an edge group of $D_{\tilde{F}}$.*

Proof. Consider the action of $F_{R(S')}$ and $\tilde{F}$ on the Bass-Serre tree T corresponding to the splitting D . Abusing notation we also use $\bar{y}$ to denote the image of $\bar{y}$ in $F_{R(S')}$.

We have hyperbolic elements $\beta, \beta' \in \tilde{F}$ such that $\beta\bar{y} = \beta'$. This means that $\bar{y}\text{Axis}(\beta') = \text{Axis}(\beta)$. Which means that there is are edges $e', e \in \text{Axis}(\beta'), \text{Axis}(\beta)$ respectively such that $\bar{y}e' = e$. On the other hand $\text{Axis}(\beta), \text{Axis}(\beta') \subset T(\tilde{F})$, the minimal $\tilde{F}$ -invariant subtree.

We also have that $D_{\tilde{F}}$ has only one edge, which means that $\tilde{F}$ is transitive on the set of edges in $T(\tilde{F})$. Let $g \in \tilde{F}$ be such that $ge = e'$. Then we have that $\bar{y}g \in \text{stab}_{\tilde{F}}(e)$. Let $\text{stab}_{F_{R(S')}}(e) \cap \tilde{F} = \langle \eta \rangle$, then we have that $\langle \eta, \bar{y}g \rangle \leq \text{stab}_{F_{R(S')}}(e)$ which is cyclic, so $\langle \eta, \bar{y}g \rangle = \langle \sqrt[n]{\eta} \rangle$. By $\tilde{F}$ -transitivity on the edges of $T(\tilde{F})$, $\text{stab}_{\tilde{F}}(e)$ for each $e \in T(\tilde{F})$ are conjugate in $\tilde{F}$ so the result follows. $\square$

It should be noted that the many ideas from the proof of Proposition 4.3 of [Sel01] are used. We now prove the lemmas.

Lemma 8.11. *Let F be a free group and let $\mu, \mu' \in F$ be two conjugate elements. Let $K = \langle \mu, \mu' \rangle \leq F$ be noncyclic and let $\gamma \in K$ be some element that is conjugate to μ in F . Then γ must be conjugate to either μ or μ' in K .*

Proof. If H is cyclic the result follows. Fix a basis of F so that μ is cyclically reduced. Suppose towards a contradiction that the result did not hold. Let $\Gamma(H)$ be the Stallings graph over F for H with basepoint v (see [KM02],[Sta83]). Let $\mu' = f^{-1}\mu f$. Then $\Gamma(H)$ is obtained by connecting two loops labelled μ by a path labelled f and folding.

Topologically, the resulting folded graph will always composed of two vertices and three arcs α, β, I . Since the graph must contain two cycles, we have two topological possibilities: either the graph can be broken into two connected components by removal of an arc; in which case we say it is *separable*; or there is no such arc.

We first consider the separable case. We have that the two cycles c_1, c_2 , as point sets, are contained in the arcs α, β . Let $u \in \Gamma(H)$ be the basepoint, w.l.o.g. it is contained in the arc α . Consider the covering space $\widetilde{\Gamma(H)}$ corresponding to the subgroup $\langle \gamma \rangle$. $\widetilde{\Gamma(H)}$ is a union of arcs that map either on to α, β or I . The core of $\widetilde{\Gamma(H)}$ must be a cycle c , moreover the cyclic word (in the basis of F) read along this arc is reduced in fact it is the cyclic word of γ . If c consists only of an arc of type α or β then we see that γ is conjugate to either μ or μ' in H . On the other hand if c contains an I -type arc, then it's length must automatically exceed that of μ , which is impossible since μ and γ are conjugate and hence must have equal cyclic words.

We now consider the non separating case. $\Gamma(H)$ consists of three arcs, moreover up to changing basis of F , we can arrange so that the path labelled μ in $\Gamma(H)$ starting at v is obtained by traversing I and then α (in particular the basepoint v is a topological vertex of $\Gamma(H)$). Since we are assuming that μ is cyclically reduced and not a proper power we have that the label of α is different from the label of I . We moreover see that, up to conjugating μ' by μ^n for some n that $(\mu')^{\pm 1}$ is the label read, starting at the basepoint v , around the cycle going first through the arc I and then through the arc β . We consider I, α, β as subwords so, abusing notation we write $\mu' = (I\beta)^{\pm 1}$ and $\mu = I\alpha$.

We have $|\alpha| = |\beta|$. Again we take the covering space $\widetilde{\Gamma(H)}$ corresponding to the subgroup $\langle \gamma \rangle$, and look at the core c . c is a cycle composed of arcs that map onto α, β, I . If c contains an arc of type I , then it must contain an arc of type either α or β . Since c has the same length as μ , c can only consist of those two arcs, which immediately implies that γ is conjugate to either μ or μ' in H . It follows that c can only consist of arcs that map onto α and β .

If $|I| \leq |\mu|/2$ then $|c| \geq |\alpha| + |\beta| > |\mu|$, which is impossible.

Suppose finally that $|I| \geq |\mu|/2$. We first make an observation about cyclically reduced words over free groups. Let $w, \tilde{w}$ be cyclically reduced words that are conjugate over the alphabet $X^{\pm 1}$ then the biinfinite words

$$\begin{array}{ccc} \dots & www & \dots \\ \dots & \tilde{w}\tilde{w}\tilde{w} & \dots \end{array}$$

viewed as maps $\mathbb{Z} \rightarrow X^{\pm 1}$ differ only by precomposition by some translation $n \mapsto n + a$ with $a < |w|$. This means that when considering the biinfinite words given by μ and μ' the subwords labelling the segment I must overlap, hence they are coherently oriented. We must therefore have

that μ' is the label of the path starting at v , going first through I and then through β . We have that γ labels a path $(\alpha^{-1}\beta)^n$ for some nonzero integer n . w.l.o.g. $\gamma = (\alpha^{-1}\beta)^n$ with $n > 0$. Since γ and μ are conjugate we must have that up to a shift the biinfinite words

$$\begin{array}{ccccccc} \dots & \alpha^{-1}\beta & \alpha^{-1}\beta & \alpha^{-1}\beta & \dots \\ \dots & & I\alpha I\alpha I\alpha & & \dots \end{array}$$

are equal. There can be no overlap between the α and α^{-1} segments, on one hand this forces $|\beta| = |I|$ which implies $|I| = |\mu|/2$. We must then have $I = \alpha^{-1}$ implying triviality of μ -contradiction. Having exhausted all possibilities the result follows. $\square$


Lemma 8.12. *If the JSJ of $\tilde{F}$ does not contain an abelian vertex group then we can always find a one edged cyclic splitting of $F_{R(S')}$ modulo F such that the induced splitting of $\tilde{F}$ also has only one edge.*

Proof. Since $\tilde{F}$ is 2 generated modulo F and is freely indecomposable modulo F , it must a group of the types previously described.

If the cyclic JSJ decomposition of $\tilde{F}$ modulo F has only one edge and one vertex, then there is nothing to show. We now consider the cyclic JSJ splittings modulo F with two edges.

If the splitting of $\tilde{F}$ has two edges and one vertex then, as we saw, either both edge classes are conjugate into F and they are conjugacy separated in F or only one of the edge classes lies in F . In both these cases only one of the edges can be visible in a one edged splitting of $F_{R(S')}$.

If the splitting of $\tilde{F}$ has two edges we have possibilities for $\tilde{F} = \mathcal{G}(A)$:




In case 2-2-C either the edges classes associated to e_1 and e_2 are both conjugate into F and are conjugacy separated or only one of them is conjugate into F , either way we can always find a one edge splitting of $F_{R(S')}$ such that only one edge is visible.

By what has been proved so far in Theorem 2.6, we know that case 2-2-A is either an amalgam of $\tilde{F}^1 \geq F$ and a free group H of rank 2 that splits as an HNN extension; or $A_v \geq F$ and A_u is abelian. In the latter case there is nothing to show. By Proposition 7.12 the edge classes associated to e_1 and e_2 are not conjugate in $\tilde{F}$, but if both edges were visible in a one edged splitting of $F_{R(S')}$ then the corresponding edge classes would be conjugate in $F_{R(S')}$. It follows that we could find a map $\phi : \tilde{F} \rightarrow F$ that is injective on H , and such that the corresponding boundary subgroups were sent to conjugates. By Lemma 7.10 and Lemma 8.11 we get that the generators of the boundary subgroups are conjugate in $\phi(H)$, and hence also in H and hence in $F_{R(S')}$ -contradiction.

In case 2-2-B we can assume that A_w is abelian so there is nothing to show.

The splittings with three edges that occur are:



In cases 2-3-B and 2-3-D, one of the vertex groups, say A_w , must be abelian so there is nothing to show.

Consider case 2-3-A. Then by the classification and Lemma 7.10, the subgroup $\bullet v \bullet \xrightarrow{e_3}$ is free of rank 2 and A_v is generated by the boundary subgroups associated to e_3 . Moreover by Proposition 7.23 we know that there must be at least two distinct edge classes. We moreover know that $A_u = F$. Consider first the case where the edge class associated to e_1 (or symmetrically e_2) is conjugate to the edge class associated to e_3 . Since the edges e_1 and e_2 are never both visible in a one-edged splitting of $F_{R(S')}$, the only possible obstruction is that, say, both e_1 and e_3 are visible in a one edged splitting of $F_{R(S')}$. If the cyclic JSJ splitting D of $F_{R(S')}$ modulo F has more than one edge then by Corollary 8.6 we can get another splitting of $F_{R(S')}$ such that only e_2 is visible. D must therefore have only one edge, and this edge must be conjugate into F . The only possibilities by Lemma 8.3 and our previous classification is that $F_{R(S')}$ is of the form $F *_u H$ where H is either free of rank 2, free abelian of rank 2 or 3, or a rank 1 extension of a centralizer with only one edge class. We look at possible induced splittings. Since the edge e_2 is not visible the subgroup $u \bullet \xrightarrow{e_2} v \bullet$ must be elliptic. Since it contains F it must be a subgroup of F which is impossible.

In case 2-3-C either there are only two edge classes, and we find ourselves as in case of 2-3-A or all edge classes are conjugate into F and F -conjugacy separated, so that only one edge will be visible in a 1 edge splitting of $F_{R(S')}$. We have exhausted all the possibilities and the result follows. $\square$

We now tackle the remaining case.

Lemma 8.13. *Suppose that the JSJ of $\tilde{F}$ has an abelian vertex group then we can find a one edge splitting of $F_{R(S')}$ modulo F such that the induced splitting of $\tilde{F}$ also has one edge.*

Proof. In this case the possible JSJs of $F_{R(S)}$ are 2-2-A, 2-2-B, 2-3-B, and 2-3-D of Lemma 8.12. Moreover obstructions only occur when the JSJ of $F_{R(S')}$ has only one edge class. In particular, if the edge class is conjugate into F , then the possibilities for $F_{R(S')}$ are

$$F *_u A, F *_u H, \text{ or } F *_u H *_u A$$

where H is free of rank 2 and A is free abelian of rank 2 or 3. We note moreover that none of the groups under consideration can be embedded into $F *_H$ since this group has no noncyclic abelian subgroups.

In case 2-2-A, Corollary 8.2 applies and we have that the edge groups associated to e_1 and e_2 cannot be conjugate.

In case 2-3-B, the vertex group $A_u = F$. The edge classes associated to e_1 and e_2 are either equal, in which case after sliding we reduce to case 2-3-D; or they are distinct in which case the edges cannot be simultaneously visible in a one edged splitting of $F_{R(S')}$. The only obstruction is that, say, the edges e_1 and e_3 are visible in a one edged splitting of $F_{R(S')}$, then we must have that the JSJ of $F_{R(S')}$ has only one edge class which forces (as in case 2-3-A of the previous proof) $F *_{A_{e_2}} A_u$ to lie in F .

In case 2-3-D we can assume that $A_u = F$. Moreover the subgroup $\bullet v \curvearrowright_{e_2}$ is free of rank 2. By Proposition 7.13 there must be at least two edge classes associated to e_1, e_2, e_3 in $\tilde{F}$. By Lemmas 7.10 and 8.11 arguing as in case 2-3-A in the previous proof, it follows that all three edges cannot be visible in a one edged splitting of $F_{R(S')}$.

Suppose first that the edges e_2 and e_3 were visible in a one edged splitting of $F_{R(S')}$, then we would have that $F *_{A_{e_1}} A_v$ is elliptic in $F_{R(S')}$. If one of the vertex groups of the JSJ of $F_{R(S')}$ is F itself, then we immediately get a contradiction. If the JSJ of $F_{R(S')}$ has two vertices and one edge, then it must be of the form

$$\tilde{F}^1 *_q H$$

where $\tilde{F}^1$ is rank 1 extension of a centralizer of F and H is free, but to get the desired induced splitting, we must have that A_w , which is free abelian of rank 2, is a subgroup of a conjugate of H which is impossible. The remaining possibility is that $F_{R(S')}$ has one edge and one vertex, but to get the induced splitting of $\tilde{F}$ in question we would need the one of the boundary subgroups of $F_{R(S')}$ to lie in a non cyclic free abelian subgroup, which by Lemma 8.3, forces $F_{R(S')} = F *_u A$ with A free abelian, and we can easily derive a contradiction.

Suppose now that only the edges e_1 and e_2 were visible in a one edge splitting of $F_{R(S')}$, then by earlier arguments we must have that the edge classes associated to e_1 and e_2 coincide in $\tilde{F}$. Then the edge class of the JSJ of $F_{R(S')}$ is conjugate into F . We also have that the subgroup $A_v *_{A_{e_3}} A_w$ is elliptic. This is clearly impossible unless $F_{R(S')} = F *_u H *_u A$ with H free and A abelian. In which case we have that $A_v *_{A_{e_3}} A_w$ is conjugate into $H *_u A$, which is only possible if u is conjugate to A_{e_3} in $F_{R(S')}$. This means that boundary subgroups associated to e_1, e_2 and e_3 are all conjugate in $F_{R(S')}$, using Lemma 8.11 and some earlier arguments we can show that the boundary subgroups associated to e_1, e_2 and e_3 are conjugate in $\tilde{F}$ —contradiction.

The remaining possibility is that only the edges e_1 and e_3 are visible in the one edged splitting. Since $\bullet v \curvearrowright_{e_2}$ is free of rank 2, we can reduce this to case 2-2-B.

We now tackle case 2-2-B. We can assume that $A_u = F$, A_v is free of rank 2, and A_w is free abelian of rank 2. We moreover have two cases the *separated* case, where $\tilde{F}$ has two edge classes; and the *unseparated* case, where $\tilde{F}$ has one edge class.

We first consider the unseparated subcase. Since the edge class is conjugate into F , any one edged splitting of $F_{R(S')}$ must have F as a

vertex group. The possibilities for $F_{R(S')}$ are

$$F *_u A \text{ or } F *_u H *_u A$$

where A is abelian and H is free. By basic Bass-Serre theory we see that $\tilde{F}$ cannot be embedded into $F *_u A$ in such a way that both edges are visible. If $F_{R(S')} = F *_u H *_u A$, then either A_v is conjugate into F or into H . If A_v is conjugate into F , then it must be embedded into some subgroup of $f a H a^{-1} f^{-1}$ with $f \in F$ and $a \in A$, then we can collapse the splitting of $F_{R(S')}$ to $(F *_u A) *_u H$ and we get a splitting of $F_{R(S')}$ modulo $\tilde{F}$ – contradiction. We therefore have that A_v is conjugable into H and A_w is conjugable into A , it follows that the splitting of $\tilde{F}$ induced by $(F *_u A) *_u H$ has only one edge.

We now consider the separated case. There are only two problematic cases for $F_{R(S')} = \pi_1(\mathcal{G}(X))$

$$\begin{aligned} \mathcal{G}(X) &= u \bullet \xrightarrow{e} \bullet v \\ \mathcal{G}(X) &= \bullet v \xleftarrow{e_1} u \bullet \xrightarrow{e_2} \bullet w \end{aligned}$$

Where $X_u = F$, X_v is noncyclic abelian and X_w is free of rank 2. Moreover all edge groups are conjugate. Consider first the case where $F_{R(S')}$ has two edges, and consider the induced splitting of $\tilde{F}$ in $F_{R(S')}$. If we collapse the edge e_2 then if the splitting of $\tilde{F}$ in this new induced splitting has only one edge then we are done, otherwise the induced splitting of $\tilde{F}$ still has two edges.

We can therefore reduce to the case where $F_{R(S')}$ is the fundamental group of

$$\mathcal{G}(X) = u \bullet \xrightarrow{e} \bullet v \quad (22)$$

with X_u non abelian and X_v noncyclic abelian. We also have $\tilde{F} = F *_p H *_q A$ which gives the graph of groups

$$\mathcal{G}(Y) = r \bullet \xrightarrow{f_1} s \bullet \xrightarrow{f_2} \bullet t \quad (23)$$

and this must be the splitting induced from $\tilde{F} \leq F_{R(S')}$ with the splitting (22). There is only one way to embed $\tilde{F}$ into $F_{R(S')}$, the embedding is

$$\tilde{F} \approx F *_p [a(\overline{H} *_p {}^g \overline{A}) a^{-1}] \quad (24)$$

where $a \in X_v$, $g \in F = X_u$, $\overline{A} \leq X_v$, and $\overline{H} \leq F$. Moreover, we have that $p, {}^g p \in \overline{H}$ but $g \notin \overline{H}$. It follows that if we construct the induced $G(X)$ -graph $\mathcal{B}$ we have

$$(F, u) \bullet \xrightarrow{(1, e, 1)} \bullet (\langle p \rangle, v) \xrightarrow{(a, e^{-1}, 1)} \bullet (\overline{H}, u) \xrightarrow{(g, e, 1)} \bullet (\overline{A}, v) \quad (25)$$

in particular the edge f_1 has length 2 and the edge f_2 has length 1. Consider the action of $\tilde{F}$ on the Bass-Serre tree T corresponding to the splitting (22) of $F_{R(S')}$.

We have elements β, β' of $\tilde{F}$ that are conjugate via the image $\bar{t}$ of the stable letter t from the splittings (19),(20) in $F_{R(S')}$. In particular we

have $\bar{t}\text{Axis}(\beta') = \text{Axis}(\beta)$. If t sends an edge $\epsilon' \in \text{Axis}(\beta')$ to an edge $\epsilon \in \text{Axis}(\beta)$ such that there is some $h \in \tilde{F}$ such that $h\epsilon' = \epsilon$ then as in the proof Lemma 8.10 we get that $F_{R(S')}$ is obtained from $\tilde{F}$ by adjoining a root to generator of an edge group, which here implies $\tilde{F} = F_{R(S')}$, and the result follows.

Otherwise let $T' = T(\tilde{F}) \leq T$ be the minimal $\tilde{F}$ -invariant subtree. Looking at (25) we see that there are three $\tilde{F}$ -orbits E_1, E_2 , and E_3 of edges in $T' \cap (\text{Axis}(\beta) \cup \text{Axis}(\beta'))$ we assume that E_2 corresponds to the edge f_2 in (23). Let $S_1, S_2, S_3 \leq \tilde{F}$ each stabilize an edge of in E_1, E_2, E_3 resp.

Consider an edge ϵ' in $\text{Axis}(\beta') \cap E_2$ that is fixed by some conjugate of the edge group $E_{f_2} \leq \tilde{F}$ of f_2 in the splitting (23). Then since $t\text{Axis}(\beta') = \text{Axis}(\beta)$. We have that tS_2t^{-1} fixes some edge say in $\epsilon \in \text{Axis}(\beta) \cap S_1$. The possibilities are limited and it follows that by conjugating boundary monomorphisms, i.e. replacing t by f_1tf_2 ; $f_1, f_2 \in \tilde{F}$, we can arrange so that $\bar{t} = aga^{-1}$, where a, g are as in (24). This means that $\bar{t}$ interchanges edges in E_1 and E_2 . It follows that $\bar{t}$ must map edges in E_3 to edges in E_3 , i.e. there is some $\epsilon \in E_3$ and some $h \in \tilde{F}$ such that $h\epsilon = \bar{t}\epsilon$ and the result follows. $\square$

We combine these two parts:

Corollary 8.14. *We can find a one edge splitting of $F_{R(S')}$ such that the induced splitting of $\tilde{F}$ also has one edge.*

8.2.2 Controlling the Strict Quotients

Proposition 8.15. *If the JSJ of $F_{R(S)}$ has only one edge and one vertex group $\tilde{F}$ that is not freely decomposable modulo F and the JSJ of $\tilde{F}$ has more than one edge and one vertex group, then $F_{R(S)}$ is an extension of a centralizer of $\tilde{F}$.*

Proof. Let $F < \tilde{F}'$ be the vertex group of the JSJ of $\tilde{F}$ that contains F . Suppose first that $\tilde{F}' = F$ or was an extension of a centralizer and that the root $\sqrt[n]{\eta}$ obtained in Lemma 8.10 was added to an edge group $\langle \eta \rangle$ lying in $\tilde{F}$. $\eta \in F$ then $\sqrt[n]{\eta} = \eta$ and $F_{R(S')} = \tilde{F}$ so the result follows. If $\tilde{F}'$ was an extension of a centralizer $F *_u \text{Ab}(u, s)$ then by our classification since either $\eta \in F$ or $\tilde{F}' = \langle F, \eta \rangle$ which means that $\eta^{\pm 1} = f_1sf_2$ with $f_i \in F - \langle u \rangle$ we have that there is no F -morphisms $F_{R(S)} \rightarrow F$ that sends η to a proper power in F so again $\sqrt[n]{\eta} = \eta$.

Otherwise we see that $F_{R(S')}$ is obtained from $\tilde{F}$ by adding elements to its vertex groups. Consider the graph of groups obtained from the JSJ of $\tilde{F}$ and adding these elements to the vertex groups. This gives a cyclic splitting D' of $F_{R(S')}$ with the same underlying graph as the JSJ of $F_{R(S)}$. It follows from our classification so far that we can in fact collapse D' to a one edge splitting so that the induced splitting of $\tilde{F}$ either has an edge group that lies in F or in $\tilde{F}$ in the way described in the previous paragraph.

It follows that $F_{R(S')} = \tilde{F}$ so $F_{R(S)}$ is an extension of a centralizer of $\tilde{F}$. $\square$

Definition 8.16. We say that two elements in $x, y \in G$ are *pseudo-conjugate* if the difference $[x] - [y]$ of their images in the abelianization of G lies in the torsion subgroup.

Proposition 8.17. *If the JSJ of $F_{R(S)}$ has only one edge and one vertex group $\tilde{F}$ which is freely indecomposable and $\pi : F_{R(S)} \rightarrow F_{R(S')}$ is a proper strict quotient then $b_1(F_{R(S')}) < b_1(F_{R(S)})$.*

Proof. If $\tilde{F}$ and $F_{R(S)}$ satisfied the hypotheses of Proposition 8.15 then the result follows immediately. We consider simultaneously the cases where the JSJ of $F_{R(S)}$ has one or two edges (i.e. (19) or (20)) replacing $\tilde{F}$ by $\bigcirc_t \alpha \tilde{F}$ if need be.

Denote by $Ab(\tilde{F})$ the abelianization of $\tilde{F}$. We see that $F_{R(S')}$ is obtained from $\tilde{F}$ by adding a the root of an element it follows that $b_1(F_{R(S')}) \leq b_1(\tilde{F})$. We consider two the elements β, β' given in (19) and (20) and consider two cases.

Suppose first that β and β' are pseudo-conjugate in $\tilde{F}$, then we see that $b_1(F_{R(S)}) > b_1(\tilde{F})$ by abelianizing a relative presentation. The result now follows.

Suppose now that β and β' are not pseudo-conjugate in $\tilde{F}$. Then in particular the element $[\beta] - [\beta']$ does not lie in the torsion subgroup. $Ab(F_{R(S')})$ can be obtained from $Ab(\tilde{F})$ by first adding the root $\sqrt[n]{\eta}$ of some element, which does not change b_1 and then identifying the images $[\beta]$ and $[\beta']$ which sends $[\beta] - [\beta']$ to zero, thus dropping the rank of the torsion free summand. The result now follows. $\square$

8.2.3 The two edge case

Proposition 8.18. *If the JSJ of $F_{R(S)}$ has two edges and one vertex then $\tilde{F}$ must be freely decomposable modulo F .*

Proof. Suppose towards a contradiction that $\tilde{F}$ was freely indecomposable modulo F . Let $F_{R(S')}$ be the first term in a strict resolution of $F_{R(S)}$ where $\tilde{F}$ splits. Then the elements β, β' must be hyperbolic elements of $F_{R(S')}$. Consider first the case where the JSJ of $\tilde{F}$ has only one edge. Then the corresponding edge group must lie in F and this splitting of $\tilde{F}$ must be induced by a one edged splitting of $F_{R(S')}$ which means that Lemma 8.10 applies, and in fact we must have that β and β' are conjugate in $\tilde{F}$, contradicting the choice of JSJ.

Suppose now that the JSJ of $\tilde{F}$ had two edges, then the possibilities are

$$F \frown H ; F \text{ --- } H \bigcirc$$

Suppose first that $F_{R(S')}$ split as an HNN extension with stable letter z and that the induced splitting of $\tilde{F}$ also had one edge. The one of the corresponding edge group either lies in F , which is impossible, or we are adding a proper root to elements of the vertex group H of $\tilde{F}$.

In this latter case, we have that β must have exponent sum 0 in z , but that β' , because $\tilde{F} = \langle F, \alpha', \beta' \rangle$ has an induced splitting as an HNN

extension, must have exponent sum 1 in z , which is impossible since they are conjugate in $F_{R(S')}$.

If $F_{R(S')}$ splits as an amalgam, and the induced splitting of $\tilde{F}$ has only one edge, then $\tilde{F}$ must also split as an amalgam and the corresponding edge group must lie in F .

If the induced splitting of $\tilde{F}$ has two edges, then either conjugation by $\bar{t}$, the image of the stable letter t , permutes the edge classes or fixes them. First note that the edge classes must be conjugacy separated in $\tilde{F}$. If conjugation by $\bar{t}$ fixes them, we can again argue that $\bar{t}$ contributes a root of an edge group lying in F , which implies that $\langle \tilde{F}, \bar{t} \rangle = \tilde{F}$.

Otherwise $\bar{t}$ permutes the edge classes which is impossible by Lemma 8.11, or by the fact that both have corresponding subgroups lying in F . $\square$

8.3 The case where $\tilde{F}$ is freely decomposable modulo F

Proposition 8.19. *Suppose that the JSJ of $F_{R(S)}$ has only one vertex group $\tilde{F}$ and suppose that $\tilde{F}$ is freely decomposable modulo F . Then the vertex group $\tilde{F}$ can only be $F * \langle z \rangle$*

Proof. By Lemmas 8.3 and 8.1 and Theorem 2.6 (A). $\tilde{F}$ must be either $F * \langle z \rangle$, $F *_u Ab(u, r) * \langle w \rangle$, $F * \langle z, w \rangle$. If $\tilde{F} = F * \langle z, w \rangle$ then Lemmas 8.3 and 8.1 would imply that $F_{R(S)}$ is freely decomposable. Applying Corollary 4.25 to the two edged case excludes $F *_u Ab(u, z) * \langle w \rangle$ as a vertex group. We now prove that $F *_u Ab(u, z) * \langle w \rangle$ cannot be a vertex group in a one edged splitting.

Suppose towards a contradiction that $F *_u Ab(u, z) * \langle w \rangle$ was the vertex group of a one edged, one vertex JSJ of $F_{R(S)}$. By Lemma 8.3 we may assume that $\tilde{F}$ is generated by $\bar{x}$ and some element β' . We know moreover that the exponent sums of z and w of words in $F \cup \{z, w\}^{\pm 1}$ representing $\bar{x}, \beta, \beta'$ do not depend on the choice of word.

$\beta \in \langle F, \bar{x} \rangle$ which means that writing β as a word in $F \cup \{\bar{x}\}^{\pm 1}$ gives equalities of exponent sums $\sigma_z(\beta) = \sigma_{\bar{x}}(\beta) * \sigma_z(\bar{x})$ and $\sigma_w(\beta) = \sigma_{\bar{x}}(\beta) * \sigma_w(\bar{x})$. Which means that we have an equality of vectors:

$$(\sigma_u(\beta), \sigma_w(\beta)) = \sigma_{\bar{x}}(\beta)(\sigma_z(\bar{x}), \sigma_w(\bar{x})) \quad (26)$$

i.e. they are linearly dependent. Now $b_1((F *_u Ab(u, z) * \langle w \rangle)) = N + 2$ which means that if $b_1(F_{R(S)}) \leq N + 2$ then β and β' must lie in the same one dimensional subgroup in the abelianization of $F \cup \{\bar{x}\}^{\pm 1}$, which means that

$$l(\sigma_z(\beta), \sigma_w(\beta)) = k(\sigma_z(\beta'), \sigma_w(\beta'))$$

for some $l, k \in \mathbb{Z}$, but this and (26) imply that $\bar{x}$ and β' cannot generate $\tilde{F}$ modulo F —contradiction. $\square$

Lemma 8.20. *Suppose the JSJ of $F_{R(S)}$ has one vertex group $\tilde{F}$ that is freely indecomposable, but does not satisfy the hypotheses of Proposition 8.15. Then the image of a strict epimorphism $F_{R(S)} \rightarrow F_{R(S')}$ must be*

$$F_{R(S')} = {}^s \left(\begin{smallmatrix} \delta \\ \delta' \end{smallmatrix} \tilde{F}' \right) \quad (27)$$

where $\tilde{F}' = F * \langle r \rangle$. Moreover $\tilde{F}$ must be of the form

$$\tilde{F} = {}_t \bigcap_{\gamma'}^{\gamma} \tilde{F}_1 \quad (28)$$

where $\tilde{F}_1 \approx F * \langle z \rangle \leq \tilde{F}'$.

Proof. By the hypotheses, we must have that the JSJ of $\tilde{F}$ has one vertex group $\tilde{F}_1$ and one edge. By Corollary 8.7 $F_{R(S')}$ must be freely indecomposable.

By Proposition 8.17, $b_1(F_{R(S')}) \leq N + 1$ where N is the rank of F . If $b_1(F_{R(S')}) = N$ then by Proposition 3.15 $F_{R(S')} = F$ which would force $\tilde{F} = F$ contradicting our hypotheses.

Let $\tilde{F} \leq \tilde{F}'$ be the vertex group of the JSJ of $F_{R(S')}$ that contains F . We have that $\tilde{F}_1 \leq \tilde{F}'$. If $\tilde{F}'$ is freely indecomposable then we could take a strict quotient of $F_{R(S')}$, apply Proposition 8.17 we would get that the image of this quotient is F , which means that $\tilde{F}' \leq F$. It would then follow that $\tilde{F}_1 = F$ and $\tilde{F}$ would have to be a rank 1 free extension of a centralizer of F . But then $\tilde{F}$ would satisfy the hypotheses of Proposition 8.15, which is a contradiction.

It therefore follows that $\tilde{F}'$ must be freely decomposable modulo F . Looking at the possibilities given in Proposition 8.19 and recalling that $b_1(F_{R(S')}) \leq N + 1$ we have that the only possibility of $\tilde{F}'$ is $\tilde{F}' \approx F * \langle r \rangle$. Since $\tilde{F}_1$ is a subgroup of $\tilde{F}'$ the only possibility is $\tilde{F}_1 \approx F * \langle z \rangle$. $\square$

Proposition 8.21. *If the JSJ of $F_{R(S)}$ has only one vertex group $\tilde{F}$, then $b_1(\tilde{F}) = N + 1$.*

Proof. In light of Proposition 8.19 and Lemma 8.20 we need only verify the case in Lemma 8.20.

Since $b_1(F_{R(S')}) \leq N + 1$ we must have that the elements δ, δ' given in (27). By Lemma 8.10, $F_{R(S')}$ is obtained from $\tilde{F}$ by the adjunction of roots of γ and γ' given in (28). $\tilde{F}_1$ is a free subgroup of the free group $\tilde{F}'$, obtained by adjoining roots. On one hand both $\tilde{F}'$ and $\tilde{F}_1$ have the same rank. On the other hand, by Theorem 7.7, $\text{Rank}(\tilde{F}') = \text{Rank}(\tilde{F}_1)$ only if γ, γ' are primitive elements of $\tilde{F}'$. Now consider the map of free abelian groups $j : A \rightarrow A'$ obtained by adding a proper root to one basis element and then adding a proper root to another basis element. Suppose that $b_1(\tilde{F}) = N + 2$. Then we must have that γ', γ are pseudo conjugate, but if that were the case, since the abelianization of $\tilde{F}$ embeds naturally into the abelianization of $\tilde{F}_1$ we must have that δ, δ' are also pseudo conjugate, which is a contradiction. $\square$

Collecting all these results gives:

Corollary 8.22. *If the JSJ of $F_{R(S)}$ has one vertex group, then all the possibilities for $F_{R(S)}$ are given by item (B) of Theorem 2.6*

References

- [App68] K. I. Appel. One-variable equations in free groups. *Proc. Amer. Math. Soc.*, 19:912–918, 1968.
- [Bau65] Gilbert Baumslag. Residual nilpotence and relations in free groups. *J. Algebra*, 2:271–282, 1965.
- [BMR99] Gilbert Baumslag, Alexei Myasnikov, and Vladimir Remeslennikov. Algebraic geometry over groups. I. Algebraic sets and ideal theory. *J. Algebra*, 219(1):16–79, 1999.
- [CR00] I. M. Chiswell and V. N. Remeslennikov. Equations in free groups with one variable. I. *J. Group Theory*, 3(4):445–466, 2000.
- [DP01] Thomas Delzant and Leonid Potyagailo. Accessibilité hiérarchique des groupes de présentation finie. *Topology*, 40(3):617–629, 2001.
- [DS99] M. J. Dunwoody and M. E. Sageev. JSJ-splittings for finitely presented groups over slender groups. *Invent. Math.*, 135(1):25–44, 1999.
- [FGM⁺98] Benjamin Fine, Anthony M. Gaglione, Alexei Myasnikov, Gerhard Rosenberger, and Dennis Spellman. A classification of fully residually free groups of rank three or less. *J. Algebra*, 200(2):571–605, 1998.
- [GK92] R. I. Grigorchuk and P. F. Kurchanov. On quadratic equations in free groups. In *Proceedings of the International Conference on Algebra, Part 1 (Novosibirsk, 1989)*, volume 131 of *Contemp. Math.*, pages 159–171, Providence, RI, 1992. Amer. Math. Soc.
- [Hme71] Ju. I. Hmelevskii. Systems of equations in a free group. I, II. *Izv. Akad. Nauk SSSR Ser. Mat.*, 35:1237–1268; *ibid.* 36 (1972), 110–179, 1971.
- [KM98a] O. Kharlampovich and A. Myasnikov. Irreducible affine varieties over a free group. I. Irreducibility of quadratic equations and Nullstellensatz. *J. Algebra*, 200(2):472–516, 1998.
- [KM98b] O. Kharlampovich and A. Myasnikov. Irreducible affine varieties over a free group. II. Systems in triangular quasi-quadratic form and description of residually free groups. *J. Algebra*, 200(2):517–570, 1998.
- [KM02] Ilya Kapovich and Alexei Myasnikov. Stallings foldings and subgroups of free groups. *J. Algebra*, 248(2):608–668, 2002.
- [KM05] Olga Kharlampovich and Alexei G. Myasnikov. Effective JSJ decompositions. In *Groups, languages, algorithms*, volume 378 of *Contemp. Math.*, pages 87–212. Amer. Math. Soc., Providence, RI, 2005.
- [KWM05] Ilya Kapovich, Richard Weidmann, and Alexei Miasnikov. Foldings, graphs of groups and the membership problem. *Internat. J. Algebra Comput.*, 15(1):95–128, 2005.

- [Lor68] A. A. Lorenc. Representations of sets of solutions of systems of equations with one unknown in a free group. *Dokl. Akad. Nauk SSSR*, 178:290–292, 1968.
- [Lys88] I. G. Lysënok. Solutions of quadratic equations in groups with the small cancellation condition. *Mat. Zametki*, 43(5):577–592, 701, 1988.
- [Mak82] G. S. Makanin. Equations in a free group. *Izv. Akad. Nauk SSSR Ser. Mat.*, 46(6):1199–1273, 1344, 1982.
- [Ol’89] A. Yu. Ol’shanskii. Diagrams of homomorphisms of surface groups. *Sibirsk. Mat. Zh.*, 30(6):150–171, 1989.
- [Ozh83] Yu. I. Ozhigov. Equations with two unknowns in a free group. *Dokl. Akad. Nauk SSSR*, 268(4):809–813, 1983.
- [Raz87] A Razborov. *On systems of equations in a free group*. PhD thesis, Steklov Math. Institute, Moscow, 1987.
- [RS97] E. Rips and Z. Sela. Cyclic splittings of finitely presented groups and the canonical JSJ decomposition. *Ann. of Math.* (2), 146(1):53–109, 1997.
- [Sel01] Zlil Sela. Diophantine geometry over groups. I. Makanin-Razborov diagrams. *Publ. Math. Inst. Hautes Études Sci.*, (93):31–105, 2001.
- [Sta83] John R. Stallings. Topology of finite graphs. *Invent. Math.*, 71(3):551–565, 1983.
- [Tou07] Nicholas W. M. Touikan. The equation $w(x,y)=u$ over free groups (preprint). <http://arxiv.org/abs/0705.4246>, 2007.
- [Wei02] Richard Weidmann. The Nielsen method for groups acting on trees. *Proc. London Math. Soc.* (3), 85(1):93–118, 2002.
- [Wic72] M. J. Wicks. A general solution of binary homogeneous equations over free groups. *Pacific J. Math.*, 41:543–561, 1972.